

Biomarkers in chronic kidney disease: a review

Robert G. Fassett^{1,2,3}, Sree K. Venuthurupalli^{3,4}, Glenda C. Gobe³, Jeff S. Coombes², Matthew A. Cooper⁵ and Wendy E. Hoy³

¹Renal Research, Royal Brisbane and Women's Hospital, Brisbane, Queensland, Australia; ²School of Human Movement Studies, The University of Queensland, Brisbane, Queensland, Australia; ³Centre for Kidney Disease Research, School of Medicine, The University of Queensland, Brisbane, Queensland, Australia; ⁴Renal Medicine, Toowoomba Hospital, Toowoomba, Queensland, Australia and ⁵Institute for Molecular Bioscience, The University of Queensland, Brisbane, Queensland, Australia

Chronic kidney disease (CKD) is a major public health problem. The classification of CKD by KDOQI and KDIGO and the routine eGFR reporting have resulted in increased identification of CKD. It is important to be able to identify those at high risk of CKD progression and its associated cardiovascular disease (CVD). Proteinuria is the most sensitive marker of CKD progression in clinical practice, especially when combined with eGFR, but these have limitations. Hence, early, more sensitive, biomarkers are required. Recently, promising biomarkers have been identified for CKD progression and its associated CVD morbidity and mortality. These may be more sensitive biomarkers of kidney function, the underlying pathophysiological processes, and/or cardiovascular risk. Although there are some common pathways to CKD progression, there are many primary causes, each with its own specific pathophysiological mechanism. Hence, a panel measuring multiple biomarkers including disease-specific biomarkers may be required. Large, longitudinal observational studies are needed to validate candidate biomarkers in a broad range of populations prior to implementation into routine CKD management. Recent renal biomarkers discovered include neutrophil gelatinase-associated lipocalin, kidney injury molecule-1, and liver-type fatty acid-binding protein. Although none are ready for use in clinical practice, it is timely to review the role of such biomarkers in predicting CKD progression and/or CVD risk in CKD.

Kidney International (2011) **80**, 806–821; doi:10.1038/ki.2011.198; published online 22 June 2011

KEYWORDS: CRP; cystatin C; neutrophil gelatinase-associated lipocalin; troponin; uric acid

Chronic kidney disease (CKD) is a major public health problem affecting 13.5% of the adult Australian¹ and 11% of the US population.² The introduction of kidney function estimating equations^{3,4} and CKD classifications by the NKF KDOQI (National Kidney Foundation Kidney Disease Outcomes Quality Initiative⁵ and the KDIGO (Kidney Disease: Improving Global Outcomes)⁶ have highlighted the condition and facilitated its diagnosis. Hence, early identification of those likely to progress to end-stage renal disease (ESRD) has become increasingly important. Existing measures such as estimated glomerular filtration rate (eGFR) and proteinuria assist with this stratification.^{7,8} However, proteinuria has limitations as a biomarker of CKD progression and response to interventions.⁹ Therefore, new validated biomarkers are required for CKD progression and cardiovascular disease (CVD) risk. Proteomic studies will provide data regarding potential biomarkers in CKD^{10–12} but these must be placed in clinical context.^{13,14} In view of recent developments in renal biomarker discovery, it is timely to review the cross-sectional and longitudinal observational studies investigating these in CKD.

This review will first address biomarkers in CKD progression and, second, biomarkers of CVD in CKD. Biomarkers are reviewed according to the main mechanisms they reflect (see Table 1 and Figure 1), although there is some overlap. Reviewed studies are summarized in Table 2. Biomarker assay variability, reliability, and stability are beyond the scope of this review.

BIOMARKER DEVELOPMENT AND TESTING IN PATIENTS

There are several clinical studies where biomarkers are being tested. The CaNPREDICT (Canadian Study of Prediction of Risk and Evolution to Dialysis, Death and Interim Cardiovascular Events Over Time) study is testing biomarkers 6 monthly in 2500 prevalent patients with eGFR 15–45 ml/min in ~50 centers across Canada over 36 months. The Spanish NEFRONA Project (Usefulness of imaging techniques and novel biomarkers in the prediction of cardiovascular risk in patients with CKD in Spain) is recruiting 2661 patients and 843 controls into a prospective observational study.¹⁵ Outcomes include cardiovascular

Correspondence: Robert G. Fassett, Level 9 Ned Hanlon Building, Royal Brisbane and Women's Hospital, Brisbane, Queensland 4029, Australia. E-mail: r.fassett@uq.edu.au

Received 1 August 2010; revised 10 January 2011; accepted 2 March 2011; published online 22 June 2011

Table 1 | Renal biomarkers according to the pathophysiological process

	Biomarker
Kidney function (GFR)	Cystatin C β-Trace protein
Tubulointerstitial injury	NGAL KIM-1 NAG L-FABP
Glomerular injury	Podocin Nephrin Podocalyxin
Endothelial dysfunction	ADMA
Oxidative stress	Ox-LDL AOPP TBARS Plasma and urinary F ₂ -isoprostanes MDA Protein reduced thiols TAS Protein carbonyls AGE Urinary 8-hydroxydeoxy guanosine 4-hydroxy-nonenal Antioxidant enzyme activities (e.g., superoxide dismutase, glutathione peroxidase, catalase) GGT
Inflammation	CRP and hs-CRP PTX3 sTNF α IL-18 Tenascin TIMP-1
Fibrosis	TGF- β 1
Cardiovascular dysfunction	ANP BNP and NT-proBNP cTnT Adrenomedullin
Metabolic disorders	Adiponectin FGF-23 ApoA-IV

Abbreviations: ADMA, asymmetric dimethylarginine; AGE, advanced glycation end product; ANP, atrial natriuretic peptide; AOPP, advanced oxidation protein products; ApoA-IV, apolipoprotein A-IV; BNP, brain natriuretic peptide; CRP, C-reactive protein; cTnT, cardiac troponin T; FGF-23, fibroblast growth factor-23; GFR, glomerular filtration rate; GGT, γ -glutamyltransferase; hs-CRP, high-sensitivity-CRP; IL-18, interleukin-18; KIM-1, kidney injury molecule-1; L-FABP, liver-type fatty acid-binding protein; MDA, malondialdehyde; NAG, *N*-acetyl- β -D-glucosaminidase; NGAL, neutrophil gelatinase-associated lipocalin; NT-proBNP, N-terminal brain natriuretic peptide; Ox-LDL, oxidized low-density lipoproteins; PTX3, pentraxin 3; sTNF α , soluble tumor necrosis factor receptor II; TAS, total antioxidant status; TBARS, thiobarbituric acid reactive substances; TGF- β 1, transforming growth factor- β 1; TIMP-1, tissue inhibitor of metalloproteinases-1.

events and mortality, carotid intima-media thickness, a composite atherosclerosis score, and biomarkers.

Other studies include the MMKD (Mild to Moderate Kidney Disease) study that recruited patients in Germany, Austria, and South Tyrol^{16,17} and reported a range of biomarkers.^{17–22} The CRIC (Chronic Renal Insufficiency Cohort)^{23,24} and the CRIB (Chronic Renal Impairment in Birmingham) study are both focusing on traditional and

nontraditional cardiovascular risk factors in CKD.²⁵ CKD studies have been reported from Japan,²⁶ China,²⁷ and Germany (Cooperative Health Research in the Region Augsburg (KORA) Study Group).²⁸ Some of these have measured multiple biomarkers.

STUDY CHARACTERISTICS AND SELECTION CRITERIA

Studies were identified through a systematic search of PubMed using key terms: biomarkers, chronic kidney disease and cardiovascular disease and papers were scrutinized for additional references.

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: KIDNEY FUNCTION

Cystatin C

Cystatin C is a small molecule (13 kDa) that is filtered and metabolized after tubular absorption.^{29,30} It is a sensitive biomarker of kidney function in mild-to-moderate kidney disease.³¹ In the MMKD study, serum cystatin C predicted CKD progression.³² The 65 of 227 patients who progressed over 7 years of follow-up had higher serum cystatin C at baseline, suggesting that it is a promising biomarker of CKD progression.³²

β-Trace protein (BTP)

BTP is a lipocalin glycoprotein^{33,34} and a more sensitive indicator of glomerular filtration than serum creatinine.^{35,36} It has been proposed as an indicator of reduced GFR, particularly in the 'creatinine-blind' range.³⁵ However, others have shown that BTP is inferior to cystatin C as an indicator of GFR.³⁷ In a report from the MMKD study group, BTP also provided reliable risk prediction for CKD progression.³² Although promising, BTP requires validation in large CKD populations.

Uric acid (UA)

Although not newly discovered, UA fulfils many criteria for a validated biomarker in CKD. It is elevated in CKD and may have a role in the pathophysiology of CKD progression through endothelial dysfunction, vascular smooth muscle cell proliferation, increased interleukin-6 (IL-6) synthesis, insulin resistance, and impairment of nitric oxide production.³⁸ Observational data indicate a relationship between serum UA and CKD prevalence and progression.^{39–41} However, not all studies have supported this contention.¹⁹ Although allopurinol lowers UA as well as improves renal function, a larger, longer duration randomized controlled trial is required.³⁸

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: KIDNEY STRUCTURE

Proteinuria

Increased urinary albumin excretion is an established predictor of CKD progression⁴² and may reflect both glomerular and tubulointerstitial injury. Analysis of data from the Nord-Trøndelag Health Study found that combining microalbuminuria with eGFR improved the ability to predict progression to ESRD.⁷ However, as a surrogate




Figure 1 | Biomarkers of chronic kidney disease (CKD) progression and renal cardiovascular disease (CVD).

marker of kidney damage, proteinuria has limitations, particularly when assessing interventions in CKD that may result in change in proteinuria.⁹ These limitations are based on the differing etiologies of CKD and the associated pathophysiological mechanisms responsible for the proteinuria in each case. In addition, the presence of proteinuria itself may cause further kidney damage.⁴³ Data from the Alberta Province laboratory registry showed that for a particular eGFR, increased proteinuria was associated with the risk of CKD progression, myocardial infarction, and mortality.⁸ Screening for microalbuminuria can detect early-stage CKD and has been shown, using microsimulation modeling, to be cost effective in patients with diabetes or hypertension.⁴⁴

Neutrophil gelatinase-associated lipocalin (NGAL)

NGAL, a ubiquitous lipocalin iron-carrying protein,⁴⁵ is highly expressed in the tubular epithelium of the distal nephron and released from tubular epithelial cells following damage such as acute kidney injury (AKI). It was originally isolated from neutrophils⁴⁶ but is also expressed in tissues such as kidney, liver, epithelial cells,^{47,48} and vascular cells in atherosclerotic plaques.⁴⁹ It has multiple molecular forms in the urine, one being dimeric originating from neutrophils and the other monomeric from kidney tubular epithelial cells.⁵⁰ This difference has the potential to improve the specificity of NGAL as a renal biomarker. Serum and

urinary NGAL have been identified as early biomarkers of AKI.^{51–58}

AKI is increasingly recognized as a prelude to CKD.⁵⁹ In an experimental study in rats, ongoing inflammation and immune activity were found to be involved with the pathogenesis of CKD, and NGAL was upregulated, suggesting that it may be a valuable biomarker for the development of CKD after AKI.^{60,61} Thus, there has been interest in NGAL as a biomarker in CKD. In addition, recent evidence suggests that NGAL may even be involved as a mediator of CKD progression.⁶² In fact, the NGAL (Lipocalin 2) knockout mouse has markedly reduced renal lesions seen in CKD progression.⁶² The same investigators in an associated retrospective clinical study in patients with autosomal dominant polycystic kidney disease found that an increase in urinary NGAL was especially seen in those patients who progressed more rapidly to end-stage kidney disease. Also, in kidney tissue from patients with IgA nephropathy and congenital nephron deficit, NGAL expression was increased in renal tubules.

In a cross-sectional study of 80 nondiabetic patients with CKD stages 2–4, serum NGAL was elevated, with the highest levels seen in those with the most advanced CKD.⁶³ Cross-sectional studies in IgA nephropathy,⁶⁴ autosomal dominant polycystic kidney disease,⁶⁵ glomerulonephritis,⁶⁶ pediatric lupus nephritis,⁶⁷ and CKD⁶⁸ have shown that urinary and serum NGAL are increased across these diseases and may

Table 2 | Studies investigating biomarkers in the progression of CKD and associated comorbidities

Authors	Biomarker	Study population	Study type	Study duration	Study outcomes
Horstrup <i>et al.</i> ⁹⁶	Tenascin and TIMP-1	Renal outpatients, Berlin, Germany (<i>n</i> =54)	Cross-sectional	NA	CKD patients had elevated serum and urinary tenascin and TIMP-1 compared with controls
Kamijo <i>et al.</i> ⁹⁰	u-L-FABP	Renal biopsy patients from St Marianna University School of Medicine (<i>n</i> =50)	Cross-sectional	NA	u-L-FABP correlated with urinary protein excretion and the degree of tubulointerstitial damage on renal biopsy
Kamijo <i>et al.</i> ⁹¹	u-L-FABP	Nondiabetic CKD patients (<i>n</i> =48)	Prospective observational	12 months	u-L-FABP was a more sensitive biomarker of CKD progression than proteinuria. u-L-FABP increased with reduced kidney function
Lee <i>et al.</i> ¹⁵⁸	GGT	CARDIA study; African-American and Caucasian males and females (<i>n</i> =2478)	Prospective observational	15 years	Serum GGT predicted microalbuminuria in those with hypertension and diabetes
Tonelli <i>et al.</i> ¹¹⁶	CRP and sTNF α	CARE study participants with CKD (<i>n</i> =687)	<i>Post hoc</i> analysis of a randomized controlled trial	58 months	Increased CRP and sTNF α at baseline predicted greater eGFR decline
Ravani <i>et al.</i> ¹⁰¹	ADMA	CKD patients, Cremona, Italy (<i>n</i> =131)	Prospective observational	7 months	ADMA levels predicted development of ESRD
Fliser <i>et al.</i> ²²	ADMA	Nondiabetic CKD, Germany, Austria, and South Tyrol (<i>n</i> =227)	Prospective observational	52.8 months	ADMA levels were significant predictors of CKD progression to ESRD and doubling of serum creatinine
Brunner <i>et al.</i> ⁶⁷	uNGAL	Pediatric SLE nephritis (<i>n</i> =35)	Cross-sectional	NA	uNGAL was significantly higher in SLE patients than controls and correlated with renal disease activity
Boes <i>et al.</i> ¹⁷	ApoA-IV	MMKD study, nondiabetic CKD (<i>n</i> =177)	Prospective observational	> 7 years	Increased ApoA-IV was associated with increased CKD progression
Mitsnefes <i>et al.</i> ⁶⁸	sNGAL	Pediatric CKD stages 2–4 (<i>n</i> =45)	Cross-sectional	NA	sNGAL correlated with cystatin C and both correlated with measured GFR and eGFR. NGAL outperformed cystatin C and eGFR at lower levels of measured GFR
Ding <i>et al.</i> ⁶⁴	uNGAL and NAG	IgA nephropathy, <i>n</i> =70 (40 Lee grade III, 30 Lee grade II)	Cross-sectional	NA	uNGAL increased in IgA Lee grade II compared with controls. uNGAL was more sensitive than NAG
Ryu <i>et al.</i> ¹⁵⁹	GGT	Korean nondiabetic, nonhypertensive males (<i>n</i> =10,337)	Prospective observational	3.5 years	Serum GGT a possible early predictor for the development of CKD
Tong <i>et al.</i> ¹²¹	PTX3	CKD stages 3 and 4 (<i>n</i> =71)	<i>Post hoc</i> analysis of a prospective observational	NA	PTX3 was negatively correlated with eGFR
Bolignano <i>et al.</i> ⁶⁵	uNGAL and sNGAL	ADPCKD (<i>n</i> =26)	Cross-sectional	NA	uNGAL and sNGAL were elevated in patients with ADPCKD compared with controls and both correlated with kidney function. Those with greater cyst development had higher uNGAL and sNGAL
Bolignano <i>et al.</i> ⁶⁶	uNGAL	Glomerulonephritis (proteinuria > 1 g/day) (<i>n</i> =33)	Cross-sectional	NA	uNGAL higher in glomerulonephritis compared with controls and significantly correlated with serum creatinine and urinary protein excretion
Lajer <i>et al.</i> ¹⁰⁰	ADMA	Type 1 diabetics with CKD (<i>n</i> =397)	Prospective observational	11.3 years	Plasma ADMA levels were associated with progression of type 1 diabetic nephropathy
Saraheimo <i>et al.</i> ¹⁴⁴	Adiponectin	Finnish Diabetic Nephropathy study, type I diabetics (<i>n</i> =1330)	Prospective observational	5 years	Adiponectin was associated with CKD progression only in those with macroproteinuria
Targher <i>et al.</i> ¹⁶⁰	GGT	NHANES (<i>n</i> =13,188)	Cross-sectional	NA	Increased GTT had a strong independent relationship with CKD
Hanai <i>et al.</i> ¹⁰²	ADMA	Japanese type 2 diabetics with normoalbuminuria or microalbuminuria (<i>n</i> =225)	Prospective observational	5.2 years	Increased ADMA levels were associated with increased risk of progression of nephropathy
Teppala <i>et al.</i> ¹⁶¹	GGT	NHANES (<i>n</i> =9516)	Cross-sectional	NA	No association between GGT and CKD
Bolignano <i>et al.</i> ⁶⁹	NGAL	White Europeans with CKD (<i>n</i> =96)	Prospective observational	18 months	Baseline urinary and serum NGAL were predictors of CKD progression
Nishida <i>et al.</i> ⁷¹	Serum and urinary NGAL	Pediatric kidney disease (<i>n</i> =85)	Prospective observational	12 months	Urinary NGAL was a superior marker of CKD in children than serum NGAL
Spanaus <i>et al.</i> ²¹	BNP and NT-proBNP	MMKD study, nondiabetic CKD (<i>n</i> =177)	Prospective observational	> 7 years	Increased BNP and NT-proBNP both predicted CKD progression
Dieplinger <i>et al.</i> ²⁰	ANP and ADM	MMKD study, nondiabetic CKD (<i>n</i> =177)	Prospective observational	> 7 years	Increased ANP and ADM both predicted CKD progression

Table 2 | Continued

Authors	Biomarker	Study population	Study type	Study duration	Study outcomes
Kiatchoosakun <i>et al.</i> ¹⁹⁸	cTnT	Stage 3 and 4 CKD (n=103)	Cross-sectional	NA	Increased cTnT was associated with reduced kidney function
Levin <i>et al.</i> ²¹²	PTH, PO ₄ , calcium	CaNPREDICT study, British Columbia, Canada, eGFR <30 ml/min per 1.73 m ² (n=4231)	Retrospective	4–31 months	CKD progression was associated with younger age, male gender, higher eGFR, higher BP, PO ₄ and PTH levels, lower Hb levels, and higher proteinuria
Sayers <i>et al.</i> ²¹³	Birth weight and other biomarkers ^a	Aboriginal Childhood Cohort Study (n=475)	Prospective cohort	11.4 years	Birth weight was negatively correlated with BP
Takamatsu <i>et al.</i> ²⁶	Urinary type IV collagen and albumin	Community study, Arita, Japan (n=1554)	Cross-sectional	NA	Albumin creatinine ratio was associated with CKD in younger participants
Fakhrzadeh <i>et al.</i> ¹¹⁸	CRP	Kahrizak Elderly Study, elderly Iranians (n=122)	Cross-sectional	NA	Combination of metabolic syndrome and high CRP strongly associated with CKD risk
Waanders <i>et al.</i> ⁷⁷	Urinary KIM-1	Proteinuric non-diabetic CKD (n=34)	Post hoc analysis of randomized controlled trial	6 weeks	Urinary KIM-1 increased in nondiabetic CKD and decreased with antiproteinuric therapy
Keller <i>et al.</i> ²¹⁴	Ten inflammatory and procoagulant biomarkers ^b	Cardiovascular Health Study (elderly) (n=4128)	Prospective cohort	7 years	Baseline serum albumin associated with rapid eGFR decline (>3 ml/min per 1.73 m ² per year) in the elderly
Fox <i>et al.</i> ¹¹⁷	CRP	Jackson Heart Study, African Americans (n=4320)	Cross-sectional	NA	CRP was associated with CKD
Manghat <i>et al.</i> ¹³⁷	FGF-23	South East Greater London, UK, Stage 1–4 CKD (n=145)	Cross-sectional	NA	FGF-23 was elevated in CKD stage 3 and correlated with diabetes, serum phosphate, and CRP
Hung <i>et al.</i> ¹¹⁹	CRP polymorphisms	AASK Study, African Americans CKD (n=642)	Prospective observational	Range 3.5–6.5 years	CRP SNPs associated with higher CRP were not associated with CKD progression. CKD patients with the rs2808630_GG genotype had a greater risk of progression
Nielsen <i>et al.</i> ⁹²	u-L-FABP	Type 1 diabetics with normoalbuminuria or microalbuminuria (n=227)	Prospective observational	18 years	High u-L-FABP predicted development and progression of diabetic nephropathy
Tsioufis <i>et al.</i> ¹⁰³	CRP and ADMA	Untreated essential hypertensives (n=13,905)	Cross-sectional	NA	Elevated CRP and ADMA were associated with microalbuminuria
Kern <i>et al.</i> ⁸⁴	Urinary NAG pentosidine, AGE fluorescence	DCCT trial type 1 diabetes (n=91)	Nested case-control	1–9 years	Baseline urinary NAG predicted macro- and micro-albuminuria. Baseline urinary AGE predicted microalbuminuria
Spanaus <i>et al.</i> ³²	Serum BTP	MMKD study, nondiabetic CKD (n=177)	Prospective observational	>7 years	BTP along with serum creatinine and cystatin C were reliable predictors of CKD progression
Wu <i>et al.</i> ⁷²	uNGAL	Drug-induced chronic tubulointerstitial nephritis (n=36)	Prospective observational	6–33 months	Baseline uNGAL was predictive of renal function decline
Kanbay <i>et al.</i> ¹⁹²	FGF-23	Gensini score in mild CKD (n=177)	Cross-sectional	NA	FGF-23 was independently associated with Gensini score, a measure of coronary atherosclerosis
Zhou <i>et al.</i> ¹³¹	Urinary CD14 mononuclear cells	Patients with ADPKD (n=16)	Prospective observational	2 years	Baseline urinary CD14 mononuclear cells showed a statistically significant correlation with kidney volume in males
Vaidya <i>et al.</i> ⁷⁹	NAG and KIM-1	Patients with type 1 diabetes mellitus (n=659)	Prospective observational	2 years	Lower urinary levels of NAG and KIM-1 were associated with regression of microalbuminuria
Viau <i>et al.</i> ⁶²	uNGAL	Patients with ADPKD (n=87)	Retrospective	6 years	Patients with higher urinary NGAL had faster progression to end-stage kidney disease

Abbreviations: AASK, African American Study of Kidney Disease and Hypertension; ADM, adrenomedullin; ADMA, asymmetric dimethylarginine; ADPKD, autosomal dominant polycystic kidney disease; AGE, advanced glycation end product; ANP, atrial natriuretic peptide; ApoA-IV, apolipoprotein A-IV; BNP, brain natriuretic peptide; BP, blood pressure; BTP, β-trace protein; CaNPREDICT study, Canadian Study of Prediction of Risk and Evolution to Dialysis, Death and Interim Cardiovascular Events Over Time study; CARDIA, Coronary Artery Risk Development in Young Adults; CARE, Cholesterol and Recurrent Events; CKD, chronic kidney disease; CRP, C-reactive protein; cTnT, cardiac troponin T; DCCT, Diabetes Control and Complications Trial; eGFR, estimated glomerular filtration rate; ESRD, end-stage renal disease; FGF-23, fibroblast growth factor-23; GGT, γ-glutamyltransferase; Hb, hemoglobin; KIM-1, kidney injury molecule-1; MMKD, Mild to Moderate Kidney Disease study; NA, not applicable; NAG, N-acetyl-β-D-glucosaminidase; NGAL, neutrophil gelatinase-associated lipocalin; NHANES, National Health and Nutrition Examination Survey; NT-proBNP, N-terminal brain natriuretic peptide; PO₄, phosphate; PTH, parathyroid hormone; PTX3, pentraxin 3; SLE, systemic lupus erythematosus; sNGAL, serum NGAL; SNP, single-nucleotide polymorphism; sTNFRII, soluble tumor necrosis factor receptor II; TIMP-1, tissue inhibitor of metalloproteinases-1; u-L-FABP, urinary liver-type fatty acid-binding protein; uNGAL, urinary NGAL.

^aBlood pressure, total cholesterol, apolipoprotein A-1, apolipoprotein B, respiratory function, ultrasound kidney size, urinary albumin/creatinine ratio, fasting triglycerides, insulin, and glucose.

^bAlbumin, CRP, interleukin-6, intercellular adhesion molecule-1, white blood cell count, fibrinogen, factor VII, factor VIII, D-dimer, and plasmin-antiplasmin complex.

reflect disease activity and kidney function. Baseline serum and urinary NGAL were predictors of eGFR decline in a longitudinal study in 96 patients with CKD from a range of etiologies over a median follow-up period of 18.5 months.⁶⁹ Both urinary and serum NGAL levels also increased in parallel in a study of diabetic patients, suggesting that NGAL may play an important role in the pathophysiology of renal adaptation to diabetes and associated nephropathy.⁷⁰ There were significant correlations between serum and urinary NGAL and eGFR and between the levels of urinary NGAL and the degree of proteinuria in 85 pediatric patients with CKD from a variety of etiologies.⁷¹ In a study of 36 patients with drug-induced chronic tubulointerstitial nephritis, increased urinary NGAL at baseline predicted decline in kidney function.⁷² There is now strong evidence that increased urinary and serum NGAL reflect damage across a spectrum of kidney diseases⁷³ as well as AKI and may predict progression of CKD.⁶⁹ The availability of more specific assays that are able to detect the different forms of NGAL may increase the specificity of urinary NGAL as a biomarker in CKD.⁵⁰

Kidney injury molecule 1 (KIM-1)

KIM-1 is a transmembrane tubular protein with uncertain function, not detectable in the normal kidney, but elevated in experimental and clinical kidney damage.⁷⁴ KIM-1 expression is significantly increased within kidney biopsy tissue collected from patients with kidney disease.⁷⁴ It is increased in the urine in AKI⁷⁵ and kidney disease,⁷⁴ and is increased in kidney transplant patients and associated with graft loss.⁷⁶ Experimental studies suggest that KIM-1 may be an indicator of AKI to CKD transition.⁶⁰ A retrospective study in proteinuric nondiabetic kidney disease patients found elevated levels of urinary KIM-1.⁷⁷ In addition, urinary KIM-1 decreased with short-term antiproteinuric therapies.⁷⁷ The diagnostic performance of KIM-1 in animal models as a predictor of drug-induced kidney injury was recently evaluated alongside more traditional serum creatinine and blood urea nitrogen and urinary *N*-acetyl- β -*D*-glucosaminidase (NAG),⁷⁸ where KIM-1 levels better correlated with the degree of kidney tubular histopathology, suggesting that KIM-1 measurement could facilitate elimination of potentially nephrotoxic drug candidates. Lower urinary levels of KIM-1 were associated with the regression of microalbuminuria in patients with type 1 diabetes mellitus.⁷⁹ Long-term observational studies in larger populations are required to assess urinary KIM-1 as a CKD biomarker.

***N*-Acetyl- β -*D*-glucosaminidase**

NAG is predominantly a biomarker of proximal tubular damage,^{80–83} but may also be a biomarker of injury to other parts of the nephron. A nested case-control study from the Diabetes Control and Compliance Trial found that baseline urinary NAG predicted both micro- and macro-albuminuria in type 1 diabetics.⁸⁴ NAG is also elevated in the urine in patients with glomerulonephritis compared with healthy

controls.⁸⁵ A recent study in type 1 diabetes mellitus found that lower levels of urinary NAG were associated with the regression of microalbuminuria.⁷⁹ It has not been assessed longitudinally in CKD.

Liver-type fatty acid-binding protein (L-FABP)

L-FABP is expressed in proximal tubular cells and is a biomarker of inflammation investigated in diabetes, diabetic nephropathy, hypertension, and early CKD.^{86–90} Urinary L-FABP (u-L-FABP) was correlated with the degree of tubulointerstitial damage and urinary protein excretion in a study of 50 patients with CKD.⁹⁰ In a prospective study, u-L-FABP was more sensitive than proteinuria in predicting CKD progression.⁹¹ Baseline u-L-FABP predicted the development of micro- and macro-albuminuria, independent of recognized biomarkers, in 165 normoalbuminuric patients from a cohort of 227 type 1 diabetics.⁹² In a recent intervention study, pitavastatin 1 mg per day reduced u-L-FABP in 30 CKD patients, which may reflect alleviation of tubulointerstitial damage.⁹³

Tenascin and tissue inhibitor of metalloproteinases 1

Tenascin is a noncollagenous matrix protein involved in interstitial renal pathology.⁹⁴ Tissue inhibitor of metalloproteinases 1 has a role in inhibition of the degradation of matrix and is upregulated in models of kidney disease.⁹⁵ Compared with healthy controls, patients with CKD have elevated serum levels of tenascin and tissue inhibitor of metalloproteinases 1 (ref. 96). Urinary levels of these molecules were also significantly raised but did not correlate with the degree of proteinuria. At this stage, these molecules are in the preliminary stages of assessment.

Glomerular injury: urinary nephrin, podocin, and podocalyxin

Although proteinuria is a feature of glomerular damage, new biomarkers of kidney podocyte injury including urinary nephrin, podocin, and podocalyxin have been discovered.^{97,98} Urinary nephrin and podocin are elevated in diabetic nephropathy and active lupus nephritis. Increased urinary podocalyxin is seen in IgA nephropathy, lupus nephritis, and post-streptococcal glomerulonephritis.⁹⁹ Although promising, and perhaps more specific for glomerular disease, these biomarkers require more detailed assessment before reaching clinical utility.

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: ENDOTHELIAL DYSFUNCTION Asymmetric dimethylarginine (ADMA)

ADMA is an amino acid that inhibits nitric oxide synthase and, if present in increased quantities, results in decreased nitric oxide production. Reduced nitric oxide production is associated with endothelial dysfunction and kidney damage. ADMA has been investigated as a biomarker in ESRD, but more recently in CKD progression.^{100–102} In a study investigating early CKD in type 1 diabetics, increased plasma levels of ADMA were predictive of the development and

progression of nephropathy.¹⁰² In another study, incident CKD patients were followed prospectively where plasma ADMA was inversely correlated with GFR and was a predictor of progression to ESRD.¹⁰¹ The progression of CKD was also greater in another population of nondiabetic CKD patients with elevated plasma ADMA levels.²² In a cross-sectional study in patients with untreated hypertension, there was an association between ADMA levels and the presence of microalbuminuria and albumin/creatinine ratio.¹⁰³

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: CARDIOVASCULAR PEPTIDES

Natriuretic peptides

Natriuretic peptides and adrenomedullin are important for cardiovascular and kidney homeostasis and their effects include natriuresis, hypotension, and diuresis.^{104,105} Natriuretic peptides have mainly been investigated as biomarkers in cardiac disease where elevated concentrations are associated with poor prognosis, degree of left ventricular dysfunction, and congestive cardiac failure.^{106–111} In CKD, N-terminal pro-brain natriuretic peptide (pro-BNP), atrial natriuretic peptide, and adrenomedullin levels have been investigated in the MMKD study where high levels of each were associated with CKD progression.^{20,21} In the first report from these investigators, BNP and N-terminal pro-BNP were measured in 177 nondiabetic CKD patients. Those patients that reached an end point, defined as doubling of serum creatinine or dialysis requirement, had significantly higher BNP and N-terminal pro-BNP levels. However, after adjustment for established predictors of CKD progression, only BNP was predictive of reaching the combined end point. Subsequently, these investigators measured plasma atrial natriuretic peptide and adrenomedullin in the same population and found that increased plasma concentrations also predicted CKD progression.²⁰ Hence, these biomarkers may prove useful in determining CKD trajectory.²¹

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: INFLAMMATION AND FIBROSIS

There are increasing numbers of inflammatory and fibrotic markers investigated in CKD and the key ones are reviewed here.

C-reactive protein (CRP) and soluble tumor necrosis factor (TNF) receptor II

CRP is a short pentraxin and an established biomarker of inflammation in kidney disease.^{112,113} TNF- α is a proinflammatory mediator, which binds to soluble TNF receptors, and plasma levels of TNF- α are increased in CKD.^{114,115} Soluble TNF receptor II is one such receptor, which is also an inflammatory marker. A *post hoc* analysis of data from the CARE (Cholesterol and Recurrent Events) trial showed that increased plasma concentrations of CRP and soluble TNF receptor II at baseline were independently associated with a greater rate of kidney function decline.¹¹⁶ A cross-sectional report from the Jackson Heart Study, a longitudinal

observational study conducted in African Americans, showed that CRP was associated with the presence of CKD.¹¹⁷ Elderly Iranians with the metabolic syndrome and high CRP levels had significantly increased risk of CKD.¹¹⁸ Single-nucleotide polymorphisms of CRP associated with higher CRP levels in African Americans were not predictive of CKD progression.¹¹⁹ However, CKD patients with the rs2808630_GG genotype had a greater risk of CKD progression. Increased CRP levels were associated with microalbuminuria in patients with untreated essential hypertension.¹⁰³

Pentraxin-3 (PTX3)

PTX3 is a long pentraxin produced in response to inflammatory signals by immune cells in contrast to CRP that is produced by hepatic cells.¹²⁰ In a study of 71 stage 3 and 4 CKD patients, there was an inverse relationship between PTX3 and eGFR.¹²¹ The investigation of this biomarker in CKD is in the very early stages.

Urinary IL-18

Urinary IL-18 is a biomarker of kidney tubular injury.¹²² In a cross-sectional study, IL-18 was markedly increased in patients with established AKI from ischemia, but not in patients with AKI from urinary tract infection, CKD, nephrotic syndrome, or those with prerenal failure.^{122,123} Urinary IL-18, more likely, offers prognostic information regarding severity and mortality at the time of AKI diagnosis, rather than indicating progression to CKD. However, evidence that patients with diabetic nephropathy have renal tubular cell overexpression of IL-18 may indicate that it deserves further study in the context of CKD.¹²⁴

Transforming growth factor- β 1 (TGF- β 1)

TGF- β 1 is a healing modulator and if secretion is increased there is accumulation of extracellular matrix and subsequent fibrosis.¹²⁵ Urinary TGF- β 1 is increased in diabetic nephropathy and glomerulonephritis.^{126,127} Serum TGF- β 1 was associated with risk factors for kidney disease in African Americans, but not Caucasians.¹²⁸ In a cross-sectional study, urine concentrations of TGF- β 1 were related to the severity of diabetic nephropathy.¹²⁹

CD14 mononuclear cells

In autosomal dominant polycystic kidney disease, where there is great heterogeneity in disease progression, GFR is known to be a poor indicator. Measurement of CD14 in the urine correlates with total kidney volume in males, and hence may prove to be a valuable early biomarker in this disease.^{130,131}

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: METABOLIC FACTORS

Fibroblast growth factor-23 (FGF-23)

FGF-23 is a hormone mediating phosphaturia and is regulated by 1,25-dihydroxyvitamin D₃. It suppresses 1 α hydroxylase activity in the proximal renal tubule.¹³² Serum

concentrations of FGF-23 are elevated in early CKD and ESRD.^{133,134} The cause of FGF-23 elevation in CKD is unclear¹³⁵ but, in early CKD, FGF-23 may rise before the rise in serum phosphate.¹³⁶ In a cross-sectional study of 145 patients with CKD stages 1–4, FGF-23 was found to be associated with other indicators of CKD complications such as diabetes, serum phosphate, and CRP. FGF-23 was elevated more in stage 3 than stage 1 and 2 CKD.¹³⁷ The phosphate binder sevelamer reduced FGF-23 levels in early-stage CKD patients with normal serum phosphate levels.¹³⁸

Apolipoprotein A-IV

Apolipoprotein A-IV is a glycoprotein involved in cholesterol transport and is increased in ESRD¹³⁹ and earlier stages of CKD.¹⁴⁰ The MMKD study demonstrated that proteinuric nondiabetic CKD patients with elevated apolipoprotein A-IV had significantly faster CKD progression.¹⁷ The elevation of apolipoprotein A-IV is reflective of kidney function and may result from reduced renal catabolism.¹⁷

Adiponectin

Adiponectin is a hormone secreted by adipocytes¹⁴¹ and is elevated in kidney disease.^{142,143} In the Finnish Diabetic Nephropathy study, adiponectin levels were only predictive of CKD progression in those with macroproteinuria, and not in those with initial normoalbuminuria or microalbuminuria.¹⁴⁴ In this context, adiponectin could not be considered a biomarker for early detection of CKD progression.

BIOMARKERS IN THE DIAGNOSIS OF CKD AND ITS PROGRESSION: OXIDATIVE STRESS

Oxidative stress refers to the *in vivo* oxidation of lipids, proteins, carbohydrates, and DNA, and it involves many biochemical pathways. Hence, there are many compounds called ‘biomarkers of oxidative stress’. Cross-sectional studies have demonstrated that CKD patients have increased levels of oxidative stress biomarkers.^{145–151} These studies have compared CKD patients with matched controls^{145,147–149} and/or compared different CKD stages.^{145–147,149,150} Biomarkers have included plasma/serum measures of oxidized low-density lipoproteins,¹⁴⁵ advanced oxidation protein products,¹⁴⁸ thiobarbituric acid reactive substances,¹⁴⁸ protein carbonyls,^{147,151} isoprostanes,^{147,150,151} protein reduced thiols,¹⁴⁷ total antioxidant status,¹⁵⁰ 4-hydroxy-nonenal,¹⁵¹ superoxide dismutase¹⁴⁹ and glutathione peroxidase¹⁴⁹ activities, urinary 8-hydroxydeoxy guanosine,¹⁴⁹ and advanced glycation end products.¹⁵¹ Advanced glycation end products are associated with the development of diabetic nephropathy.¹⁵² Pentosidine is an advanced glycation end product, which may be present in the urine.¹⁵³ In a nested case-control study from the Diabetes Control and Compliance Trial, baseline urinary pentosidine predicted macroalbuminuria and baseline advanced glycation end product predicted microalbuminuria.⁸⁴

Findings from studies in CKD patients include: (1) oxidized low-density lipoproteins are associated with endothelial

injury¹⁴⁵ and inflammation,¹⁴⁸ (2) elevated oxidative stress occurs early in CKD,¹⁴⁷ and increases as CKD progresses,¹⁵⁰ and (3) patients with diabetic nephropathy^{147,151} and proteinuria¹⁴⁹ have additional elevations of oxidative stress. Oxidative stress increases further on renal replacement therapy^{145,146} and regresses after kidney transplantation.¹⁵⁴

Of most relevance to this review is the equivocal relationship between eGFR and biomarkers of oxidative stress. Two studies reported that oxidative stress was related to kidney function,^{145,150} and one reported no link.¹⁴⁷ This difference may relate to the biomarkers measured and highlights the complexity of oxidative stress assessment.

Future research should include the direct assessment of reactant generation at the tissue level, use of multiple biomarkers for a more comprehensive assessment of pathways involved, and longitudinal cohort studies assessing how changes in these measures are related to outcomes such as morbidity and mortality.

γ-Glutamyl transpeptidase (GGT)

Until recently, the epidemiological associations of GGT were unexplored.¹⁵⁵ GGT has mainly been used as a marker of liver disease and alcohol intake,¹⁵⁶ but it has also been touted as a biomarker of oxidative stress.¹⁵⁷ In the CARDIA (Coronary Artery Risk Development in Young Adults) study, those with hypertension and diabetes had an association between GGT (within the normal range) and the occurrence of microalbuminuria.¹⁵⁸ Ryu *et al.*¹⁵⁹ reported, in a cohort of nonhypertensive and nondiabetic Korean males, that GGT may be an early predictor of CKD even when allowing for confounding factors at baseline. In a cross-sectional analysis of 13,188 adults in the NHANES (National Health and Nutrition Examination Survey) 2001–2006, there was a strong independent association between elevated GGT and CKD.¹⁶⁰ Adjustment was undertaken for comorbidities, demographics, alcohol consumption, lipid-lowering therapy, the presence of viral hepatitis, and laboratory results. However, Teppala *et al.*¹⁶¹ reported cross-sectional data from NHANES 1999–2002 showing no association between increased GGT levels and CKD when adjusted for age, gender, race, smoking, alcohol consumption, diabetes, hypertension, obesity, and serum lipids. The reasons for the differences between these studies are unclear.

DISEASE-SPECIFIC BIOMARKERS

Although elements of CKD progression are common to all underlying primary renal etiologies, there are specific pathophysiological mechanisms linked to the primary renal diseases underpinning the diagnosis of CKD. For this reason, it is unlikely that a single biomarker will be able to predict CKD progression when encompassing all primary renal diseases. Disease-specific markers may supplement more general biomarkers.

For instance, in idiopathic membranous nephropathy, 70% of patients from one study had the M-type phospholipase A₂ receptor identified in glomerular extracts, and

the presence of antibodies against this antigen supported this as important in the disease.¹⁶² In addition- single nucleotide polymorphisms of the phospholipase A₂ receptor 1 may be involved in the etiology of idiopathic membranous nephropathy.^{163,164}

BIOMARKERS OF CVD MORBIDITY AND MORTALITY IN CKD

CVD is intimately associated with CKD and its progression, and hence it is not surprising that there has been a keen interest in searching for biomarkers that might predict CVD morbidity and mortality in the CKD population. Along with traditional risk factors, nontraditional risk factors contribute to CVD in CKD. The next section reviews biomarkers that have been investigated in the context of predicting CVD morbidity and mortality in CKD. These studies are summarized in Table 3.

Cystatin C

As plasma cystatin C is an accurate biomarker of kidney function, and the latter is closely correlated with CVD risk, it follows that plasma cystatin C would also be a biomarker of CVD risk. Prospective studies have demonstrated an increased CVD risk associated with increased plasma

cystatin C levels.¹⁶⁵ In fact, in elderly individuals without CKD, plasma cystatin C is a biomarker for the risk for CVD and mortality^{166,167} and reflects coronary artery disease severity.¹⁶⁸ In patients with CKD, elevated plasma levels of cystatin C are associated with all-cause mortality, cardiovascular events, and incident heart failure.¹⁶⁹ Plasma cystatin C is, therefore, a powerful predictor of CVD and mortality in those with and without CKD.

Uric acid

Although there are some observational data linking serum UA with cardiovascular and all-cause mortality in the general population, there is little information assessing this relationship in CKD.¹⁷⁰ Analysis of data from the ARIC (Atherosclerosis Risk In Communities) study found that although there was an independent association between elevated serum UA and mortality, there were no significant associations for cardiovascular events and mortality in the CKD population.¹⁷⁰

Natriuretic peptides

BNP was prospectively studied as a biomarker of cardiovascular events in a CKD population in Japan, where the relative risk of cardiovascular events was significantly higher in those

Table 3 | Studies investigating biomarkers in CVD morbidity and mortality in CKD

Authors	Biomarker	Study population	Study type	Study duration	Study outcomes
Tong <i>et al.</i> ¹²¹	PTX3	CKD stages 3 and 4 (n=71)	Post hoc analysis of prospective cohort	NA	PTX3 was positively correlated with cardiovascular events and mortality
Goicoechea <i>et al.</i> ¹⁸⁴	Fibrinogen	CKD patients with eGFR <60 ml/min per 1.73 m ² (n=128)	Prospective observational longitudinal	5.5 years	Serum fibrinogen predicted all-cause mortality in CKD stages 3-5
Weiner <i>et al.</i> ¹⁸⁵	Fibrinogen, albumin, triglyceride, and CRP	ARIC study participants, eGFR 60 ml/min per 1.73 m ² (n=1678)	Prospective cohort study	108 months	Increased fibrinogen, triglyceride, CRP, decreased albumin, predicted composite events (MI, stroke, all-cause mortality)
Levin <i>et al.</i> ²¹²	PTH, PO ₄ , calcium	CaNPREDICT study, British Columbia, Canada, eGFR <30 ml/min per 1.73 m ² (n=4231)	Retrospective	4-31 months	Death associated with older age, lower BP, Hb, higher PO ₄ , PTH. Vitamin D associated with reduced death
Lajer <i>et al.</i> ¹⁰⁰	ADMA	Type 1 diabetics with CKD (n=397)	Prospective observational	11.3 years	Plasma ADMA levels predicted fatal and nonfatal cardiovascular events
Ravani <i>et al.</i> ¹⁰¹	ADMA	CKD patients, Cremona, Italy (n=131)	Prospective observational study	27 months	ADMA levels predicted mortality
Liabeuf <i>et al.</i> ¹⁹⁶	sTRAIL	CKD stages 2-5 included 31% CKD stage 5D (n=130)	Prospective cohort	20 months-2.1 years	sTRAIL inversely associated with mortality
Junyent <i>et al.</i> ¹⁵	Multiple biomarkers	NEFRONA Project Spain (n=2661)	Prospective cohort (in progress)	4 years	Cardiovascular events, mortality, surrogate vascular measures, and biomarkers (in progress)
Sakuma <i>et al.</i> ¹⁷¹	BNP	Iwate-KENCO study, Ninohe, Japan (n=15,394)	Prospective observational	2.8 years	BNP a strong predictor of cardiovascular events
Nielsen <i>et al.</i> ⁹²	u-L-FABP	Type 1 diabetics with normoalbuminuria or microalbuminuria (n=227)	Prospective observational	18 years	High u-L-FABP predicted mortality
Seiler <i>et al.</i> ¹⁹¹	FGF-23	CKD patients not on dialysis (n=149)	Prospective observational	4.8 years	Elevated FGF-23 were predictive of cardiovascular events
Rogacev <i>et al.</i> ²⁰⁷	CD14++ and CD16+ monocytes	CKD patients not on dialysis (n=119)	Prospective observational	4.9 years	Presence of CD14++ and CD16+ monocytes were independently associated with cardiovascular events

Abbreviations: ADMA, asymmetric dimethylarginine; ANP, atrial natriuretic peptide; ARIC, Atherosclerosis Risk In Communities; BP, blood pressure; BNP, brain natriuretic peptide; CaNPREDICT study, Canadian Study of Prediction of Risk and Evolution to Dialysis, Death and Interim Cardiovascular Events Over Time; CARDIA, Coronary Artery Risk Development in Young Adults; CKD, chronic kidney disease; CRP, C-reactive protein; cTnT, cardiac troponin T; CVD, cardiovascular disease; eGFR, estimated glomerular filtration rate; FGF-23, fibroblast growth factor-23; GGT, γ -glutamyltransferase; Hb, hemoglobin; MI, myocardial infarction; NA, not applicable; NHANES, National Health and Nutrition Examination Survey; NT-proBNP, N-terminal brain natriuretic peptide; PO₄, phosphate; PTH, parathyroid hormone; PTX3, pentraxin 3; sTRAIL, soluble tumor necrosis factor-related apoptosis-inducing ligand; u-L-FABP, urinary liver-type fatty acid-binding protein.

with the highest BNP levels.¹⁷¹ In children with CKD, increased levels of BNP were associated with ventricular hypertrophy with higher levels present in eccentric than concentric hypertrophy, suggesting that BNP may be predictive of abnormal cardiac geometry.¹⁷² The natriuretic peptides have proven useful CVD biomarkers in the general population and have potential for similar use in CKD patients pending further studies.

Asymmetric dimethylarginine

ADMA is a biomarker of atherosclerosis¹⁷³ and is associated with increased blood pressure¹⁷⁴ and insulin resistance.¹⁷⁵ It is a strong independent predictor of cardiovascular and mortality risk in end-stage kidney disease,¹⁷⁶ but has not been as extensively investigated in predialysis CKD. In a prospective study in 131 incident CKD patients followed for a mean of 27 months, plasma ADMA levels were independently predictive of total mortality.¹⁰¹ Also, in another prospective observational study of type 1 diabetics with overt nephropathy, plasma ADMA was predictive of both fatal and nonfatal cardiovascular events.¹⁰⁰ Further studies of this biomarker in more diverse CKD populations are required.

Fibrinogen

Serum fibrinogen is a predictor of CVD events in the general population,^{177–180} and is independently predictive of cardiovascular and all-cause mortality in end-stage kidney disease.¹⁸¹ This association has been demonstrated in patients with CKD.^{182,183} More recent studies in patients with stage 3 and 4 CKD indicated that an elevated serum fibrinogen level was an independent predictor of all cause mortality.¹⁸⁴ Weiner *et al.*¹⁸⁵ assessed nontraditional cardiovascular risk factors including serum fibrinogen in stage 3 and 4 CKD patients in both the ARIC and Cardiovascular Health studies where increased serum fibrinogen was independently predictive of composite events that included myocardial infarction, stroke, and all-cause mortality. Serum fibrinogen may prove useful in a cardiovascular biomarker panel for identifying CKD patients at risk of CVD.

Neutrophil gelatinase-associated lipocalin

Recent interest has focused on NGAL as a biomarker of CVD as well as in AKI and CKD progression.¹⁸⁶ This stems from experimental¹⁸⁷ and clinical evidence in the nonrenal population.^{188,189} However, in a recent pilot study of 46 elderly patients with heart failure and eGFR 72.3 ± 15.1 ml/min per 1.73 m^2 , plasma NGAL levels increased in parallel with predictive indicators of CVD, with plasma NGAL levels > 783 ng/ml correlating with CVD mortality after 2 years.¹⁹⁰ NGAL may prove to be a useful biomarker of CVD associated with CKD but studies in CKD populations are required.

Fibroblast growth factor-23

Elevated levels of FGF-23 have been shown to be predictive of mortality in ESRD and cardiovascular events in CKD patients before starting dialysis.¹⁹¹ A cross-sectional study in patients

with mild CKD found an independent association between increased FGF-23 and the Gensini score, a measure of total coronary atherosclerosis.¹⁹² Additional validation of this association is required.¹⁹³

Pentraxin-3

Only one study assessing PTX3 as a biomarker of CVD in CKD was identified, where PTX3 was associated with protein-energy wasting, inflammation, CVD, and all-cause mortality in a study of 71 patients with stage 3 and 4 CKD.¹²¹

Tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)

TRAIL is a factor associated with protection against inflammation and is involved with apoptosis¹⁹⁴ and atherosclerosis.¹⁹⁵ Liabeuf *et al.*¹⁹⁶ assessed soluble TRAIL in 130 patients with stage 2–5 CKD that included 31% ESRD patients. Soluble TRAIL was found to be a biomarker associated with reduced inflammation and mortality.

Troponin T and I

Troponin T and I are sensitive biomarkers of cardiac injury in the general population. Although mechanisms vary for causation, increased plasma Troponin T and I, above the cutoffs for a normal population, were independently predictive of subsequent death in ESRD patients.¹⁹⁷ Elevations in troponin T (cTnT) were frequent findings in patients with stage 3 and 4 CKD and elevated cTnT has been associated with reduced renal clearance.¹⁹⁸ Elevated cTnT in patients with CKD has also been shown to be because of reduced renal clearance in patients with congestive heart failure.¹⁹⁹ However, detectable cTnT is associated with reduced survival.²⁰⁰ The same investigators found cTnT and cTnI elevated in CKD and that elevated cTnI was associated with increased mortality.²⁰¹ In addition, elevated levels of cTnT and cTnI are seen frequently in patients with CKD predialysis even in the absence of an acute coronary syndrome. In fact, it has been shown that elevated levels of cTnT measured in patients before starting dialysis predict asymptomatic multivessel coronary artery stenosis.²⁰² cTnT and cTnI were found to be equivalent in differentiating short-term prognosis in patients with CKD with non-ST-segment elevation acute coronary syndromes but this was more useful in those with more advanced CKD.²⁰³ Thus, despite elevations in these biomarkers of CVD being seen associated with reduced renal clearance, they are of use as biomarkers of mortality risk in predialysis CKD patients.

Liver-type fatty acid-binding protein

In a prospective cohort study of 165 normoalbuminuric type 1 diabetics, u-L-FABP was predictive of all-cause mortality, independent of traditional risk factors such as gender, age, smoking status, HbA_{1c}, blood pressure, urinary albumin excretion rate, and kidney function.⁹² Although this study was not performed in the CKD population, it explored progression to proteinuria and u-L-FABP may be worth

assessing as a candidate biomarker of CVD and mortality in CKD patients.

CD14++ CD16+ monocytes

Macrophages and monocytes are recognized participants in the atherosclerotic process.²⁰⁴ Subsets of monocytes have been identified based on the expression of different receptors (lipopolysaccharide CD14 and FcγIII CD16) on their surface. High counts of CD14++ CD16+ monocytes are increased in inflammation²⁰⁵ and are also associated with the development of cardiovascular events in the dialysis setting.²⁰⁶ In a prospective study of 119 patients with predialysis CKD from a range of etiologies, the presence of CD14++ and CD16+ monocytes was associated independently with cardiovascular events.²⁰⁷ At this stage, these results are purely a stimulus for further investigation.

NEW BIOMARKER DEVELOPMENT

New biomarkers are usually identified from experimental studies before investigation in clinical populations. Recently, studies have investigated predictive biomarkers of acute drug-induced kidney injury.^{78,208–210} For example, in a rodent study of nephrotoxin-induced injury, urinary TFF3 (trefoil factor 3) protein levels were markedly reduced, and urinary albumin increased in association with tubular injury. *In situ* hybridization of TFF3 mRNA also decreased in injured tubules. Albumin outperformed serum creatinine and blood urea nitrogen for detecting tubular injury, but TFF3 augmented the potential of blood urea nitrogen and serum creatinine to detect kidney damage.²¹⁰ The Nephrotoxicity Working Group within the Critical Path Institute, a pharmaceutical industry public-private partnership, submitted a recommendation to the US FDA (US Food and Drug Administration) and the European Medicines Agency for further study of seven urinary biomarkers of kidney injury (KIM-1, albumin, total protein, β₂-microglobulin, cystatin C, clusterin, and trefoil factor 3), which qualified for use in regulatory decision-making (Table 4). Most biomar-

kers in the panel showed better sensitivity and specificity than blood urea nitrogen and serum creatinine, and all added complementary information.²¹¹ Testing those not currently assessed in CKD populations may prove fruitful. The sequence of biomarker discovery and development through to clinical utility is summarized in Figure 2.

SUMMARY

To enable nephrologists to focus on CKD patients that will have a poor trajectory, we need better biomarkers that can identify this earlier. Serum creatinine, eGFR, and proteinuria are insensitive and reliance on these may result in extensive time lapse where successful interventions could be tested and applied. Some biomarkers reviewed show promise but further validation is required in larger, more diverse populations before translation into clinical practice. Of those




Figure 2 | Summary of biomarker discovery and development through to clinical utility.

Table 4 | Summary of claims submitted to the FDA and EMEA on biomarkers associated with nephrotoxicity

Biomarker	Qualified preclinically	Adds information to SCr and BUN ^{a,b}	Outperforms SCr and/or BUN ^{a,b,c,d}	Analytically validated assay	Widely available assay	Qualified for clinical use ^e
KIM-1	Yes	Yes ^a	Yes ^a	Yes	Pending	Yes
Albumin	Yes	Yes ^a	Yes ^a	Yes	Yes	Yes
CLU	Yes	Yes ^a	Yes ^a	Yes	Yes	Pending
TFF3	Yes	Yes ^a	No	Yes	Pending	Pending
Total protein	Yes	Yes ^b	Yes ^{b,c}	Yes	Yes	Yes
Cystatin C	Yes	Yes ^b	Yes ^{b,c}	Yes	Yes	Yes
β ₂ -Microglobulin	Yes	Yes ^b	Yes ^{b,c}	Yes	Yes	Yes

Abbreviations: BUN, blood urea nitrogen; CLU, clusterin; EMEA, European Medicines Agency; FDA, Food and Drug Administration; KIM-1, kidney injury molecule-1; SCr, serum creatinine; TFF3, trefoil factor 3.

^aAcute tubular alterations.

^bAcute glomerular injury with acute tubular reabsorption impairment.

^cBiomarker outperformed SCr.

^dIf an inclusion receiver operating characteristic (ROC) analysis (true positive rate (sensitivity) against the rate of false positives (1-specificity) for a continuous variable against a specific reference standard) analysis is considered, instead of an exclusion ROC analysis, cystatin C and β₂-microglobulin outperform not only SCr but also BUN with respect to the prediction of histopathologically confirmed kidney injury (see Honkanen *et al.*¹²⁶ for further details of ROC analysis).

^eQualified for clinical use refers to a 'case-by-case' on text and not to a broad general qualification.

reviewed, NGAL has the greatest promise as a biomarker of CKD progression and cystatin C as a biomarker of kidney function, CKD progression, and cardiovascular risk. It is unlikely that a single marker will satisfy the requirement of predicting CKD progression and cardiovascular morbidity and mortality as it would be unlikely to reflect the complexities of the multiple pathophysiological processes involved in CKD progression or the underlying primary renal disease. It is more likely that a focused panel of biomarkers will be most rewarding.

In conclusion, the search for new relevant biomarkers to better stratify patients with CKD according to the risk of progression, morbidity, and mortality is underway. It is important to determine whether the newly identified biomarkers are purely associations or real biomarkers of underlying pathophysiological processes. Testing biomarkers prospectively in large, divergent populations over extended follow-up periods, and validating them against hard outcome measures such as the development of ESRD and mortality is required before translation into clinical practice. Although advances in proteomics technologies, sample conditioning, and analysis methods have greatly improved productivity and efficiency in biomarker discovery, biomarker verification and validation remains a significant, costly, and high-risk undertaking in the commercial development and deployment of novel biomarkers for CKD.

DISCLOSURE

All the authors declared no competing interests.

ACKNOWLEDGMENTS

The Centre for Chronic Disease at the University of Queensland has funded this work.

AUTHOR CONTRIBUTIONS

RGF, SKV, GCG, JSC, MAC, and WEH are responsible for writing the manuscript. All authors have read and approved the final manuscript.

REFERENCES

- Chadban SJ, Briganti EM, Kerr PG *et al.* Prevalence of kidney damage in Australian adults: the AusDiab kidney study. *J Am Soc Nephrol* 2003; **14**(7 Suppl 2): S131–S138.
- Coresh J, Astor BC, Greene T *et al.* Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2003; **41**: 1–12.
- Levey AS, Stevens LA, Schmid CH *et al.* A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009; **150**: 604–612.
- Levey AS, Coresh J, Greene T *et al.* Expressing the Modification of Diet in Renal Disease Study equation for estimating glomerular filtration rate with standardized serum creatinine values. *Clin Chem* 2007; **53**: 766–772.
- National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* 2002; **39**(2 Suppl 1): S1–S266.
- Levey AS, Atkins R, Coresh J *et al.* Chronic kidney disease as a global public health problem: approaches and initiatives—a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 2007; **72**: 247–259.
- Hallan SI, Ritz E, Lydersen S *et al.* Combining GFR and albuminuria to classify CKD improves prediction of ESRD. *J Am Soc Nephrol* 2009; **20**: 1069–1077.
- Hemmelgarn BR, Zhang J, Manns BJ *et al.* Nephrology visits and health care resource use before and after reporting estimated glomerular filtration rate. *JAMA* 2010; **303**: 1151–1158.
- Levey AS, Catran D, Friedman A *et al.* Proteinuria as a surrogate outcome in CKD: report of a scientific workshop sponsored by the National Kidney Foundation and the US Food and Drug Administration. *Am J Kidney Dis* 2009; **54**: 205–226.
- Ben Ameer R, Molina L, Bolvin C *et al.* Proteomic approaches for discovering biomarkers of diabetic nephropathy. *Nephrol Dial Transplant* 2010; **25**: 2866–2875.
- Smith MP, Banks RE, Wood SL *et al.* Application of proteomic analysis to the study of renal diseases. *Nat Rev Nephrol* 2009; **5**: 701–712.
- Ben Ameer R, Molina L, Bolvin C *et al.* Proteomic approaches for discovering biomarkers of diabetic nephropathy. *Nephrol Dial Transplant* 2010; **25**: 2866–2875.
- Knepper MA. Common sense approaches to urinary biomarker study design. *J Am Soc Nephrol* 2009; **20**: 1175–1178.
- Neilson EG. Finding new sea legs for urine proteomics. *J Am Soc Nephrol* 2009; **20**: 1162.
- Junyent M, Martinez M, Borrás M *et al.* [Usefulness of imaging techniques and novel biomarkers in the prediction of cardiovascular risk in patients with chronic kidney disease in Spain: the NEFRONA project]. *Nefrologia* 2010; **30**: 119–126.
- Kronenberg F, Kuen E, Ritz E *et al.* Lipoprotein(a) serum concentrations and apolipoprotein(a) phenotypes in mild and moderate renal failure. *J Am Soc Nephrol* 2000; **11**: 105–115.
- Boes E, Fliser D, Ritz E *et al.* Apolipoprotein A-IV predicts progression of chronic kidney disease: the mild to moderate kidney disease study. *J Am Soc Nephrol* 2006; **17**: 528–536.
- Fliser D, Kollerits B, Neyer U *et al.* Fibroblast growth factor 23 (FGF23) predicts progression of chronic kidney disease: the Mild to Moderate Kidney Disease (MMKD) Study. *J Am Soc Nephrol* 2007; **18**: 2600–2608.
- Sturm G, Kollerits B, Neyer U *et al.* Uric acid as a risk factor for progression of non-diabetic chronic kidney disease? The Mild to Moderate Kidney Disease (MMKD) Study. *Exp Gerontol* 2008; **43**: 347–352.
- Dieplinger B, Mueller T, Kollerits B *et al.* Pro-A-type natriuretic peptide and pro-adrenomedullin predict progression of chronic kidney disease: the MMKD Study. *Kidney Int* 2009; **75**: 408–414.
- Spanaus KS, Kronenberg F, Ritz E *et al.* B-type natriuretic peptide concentrations predict the progression of nondiabetic chronic kidney disease: the Mild-to-Moderate Kidney Disease Study. *Clin Chem* 2007; **53**: 1264–1272.
- Fliser D, Kronenberg F, Kielstein JT *et al.* Asymmetric dimethylarginine and progression of chronic kidney disease: the mild to moderate kidney disease study. *J Am Soc Nephrol* 2005; **16**: 2456–2461.
- Feldman HI, Appel LJ, Chertow GM *et al.* The Chronic Renal Insufficiency Cohort (CRIC) Study: design and methods. *J Am Soc Nephrol* 2003; **14**(7 Suppl 2): S148–S153.
- Lash JP, Go AS, Appel LJ *et al.* Chronic Renal Insufficiency Cohort (CRIC) Study: baseline characteristics and associations with kidney function. *Clin J Am Soc Nephrol* 2009; **4**: 1302–1311.
- Wheeler DC, Townsend JN, Landray MJ. Cardiovascular risk factors in predialysis patients: baseline data from the Chronic Renal Impairment in Birmingham (CRIB) study. *Kidney Int Suppl* 2003; **84**: S201–S203.
- Takamatsu N, Abe H, Tominaga T *et al.* Risk factors for chronic kidney disease in Japan: a community-based study. *BMC Nephrol* 2009; **10**: 34.
- Shan Y, Zhang Q, Liu Z *et al.* Prevalence and risk factors associated with chronic kidney disease in adults over 40 years: a population study from Central China. *Nephrology (Carlton)* 2010; **15**: 354–361.
- Baumeister SE, Boger CA, Kramer BK *et al.* Effect of chronic kidney disease and comorbid conditions on health care costs: a 10-year observational study in a general population. *Am J Nephrol* 2010; **31**: 222–229.
- Randers E, Erlandsen EJ. Serum cystatin C as an endogenous marker of the renal function—a review. *Clin Chem Lab Med* 1999; **37**: 389–395.
- Madero M, Sarnak MJ, Stevens LA. Serum cystatin C as a marker of glomerular filtration rate. *Curr Opin Nephrol Hypertens* 2006; **15**: 610–616.
- Newman DJ, Thakkar H, Edwards RG *et al.* Serum cystatin C measured by automated immunoassay: a more sensitive marker of changes in GFR than serum creatinine. *Kidney Int* 1995; **47**: 312–318.
- Spanaus KS, Kollerits B, Ritz E *et al.* Serum creatinine, cystatin C, and beta-trace protein in diagnostic staging and predicting progression of primary nondiabetic chronic kidney disease. *Clin Chem* 2010; **56**: 740–749.
- Hoffmann A, Nimtz M, Conradt HS. Molecular characterization of beta-trace protein in human serum and urine: a potential diagnostic marker for renal diseases. *Glycobiology* 1997; **7**: 499–506.

34. Melegos DN, Grass L, Pierratos A *et al.* Highly elevated levels of prostaglandin D synthase in the serum of patients with renal failure. *Urology* 1999; **53**: 32–37.
35. Priem F, Althaus H, Birnbaum M *et al.* Beta-trace protein in serum: a new marker of glomerular filtration rate in the creatinine-blind range. *Clin Chem* 1999; **45**: 567–568.
36. Dharmidharka VR, Kwon C, Stevens G. Serum cystatin C is superior to serum creatinine as a marker of kidney function: a meta-analysis. *Am J Kidney Dis* 2002; **40**: 221–226.
37. Woitas RP, Stoffel-Wagner B, Poege U *et al.* Low-molecular weight proteins as markers for glomerular filtration rate. *Clin Chem* 2001; **47**: 2179–2180.
38. Siu YP, Leung KT, Tong MK *et al.* Use of allopurinol in slowing the progression of renal disease through its ability to lower serum uric acid level. *Am J Kidney Dis* 2006; **47**: 51–59.
39. Chonchol M, Shlipak MG, Katz R *et al.* Relationship of uric acid with progression of kidney disease. *Am J Kidney Dis* 2007; **50**: 239–247.
40. Chang HY, Tung CW, Lee PH *et al.* Hyperuricemia as an independent risk factor of chronic kidney disease in middle-aged and elderly population. *Am J Med Sci* 2010; **339**: 509–515.
41. Feig DI. Uric acid: a novel mediator and marker of risk in chronic kidney disease? *Curr Opin Nephrol Hypertens* 2009; **18**: 526–530.
42. Taal MW, Brenner BM. Renal risk scores: progress and prospects. *Kidney Int* 2008; **73**: 1216–1219.
43. Remuzzi G, Benigni A, Remuzzi A. Mechanisms of progression and regression of renal lesions of chronic nephropathies and diabetes. *J Clin Invest* 2006; **116**: 288–296.
44. Hoerger TJ, Wittenborn JS, Segel JE *et al.* A health policy model of CKD: 2. The cost-effectiveness of microalbuminuria screening. *Am J Kidney Dis* 2010; **55**: 463–473.
45. Bolignano D, Coppolino G, Romeo A *et al.* Neutrophil gelatinase-associated lipocalin (NGAL) reflects iron status in haemodialysis patients. *Nephrol Dial Transplant* 2009; **24**: 3398–3403.
46. Kjeldsen L, Johnsen AH, Sengelov H *et al.* Isolation and primary structure of NGAL, a novel protein associated with human neutrophil gelatinase. *J Biol Chem* 1993; **268**: 10425–10432.
47. Cowland JB, Borregaard N. Molecular characterization and pattern of tissue expression of the gene for neutrophil gelatinase-associated lipocalin from humans. *Genomics* 1997; **45**: 17–23.
48. Friedl A, Stoesz SP, Buckley P *et al.* Neutrophil gelatinase-associated lipocalin in normal and neoplastic human tissues. Cell type-specific pattern of expression. *Histochem J* 1999; **31**: 433–441.
49. Hemdahl AL, Gabrielsen A, Zhu C *et al.* Expression of neutrophil gelatinase-associated lipocalin in atherosclerosis and myocardial infarction. *Arterioscler Thromb Vasc Biol* 2006; **26**: 136–142.
50. Cai L, Rubin J, Han W *et al.* The origin of multiple molecular forms in urine of HNL/NGAL. *Clin J Am Soc Nephrol* 2010; **5**: 2229–2235.
51. Mishra J, Dent C, Tarabishi R *et al.* Neutrophil gelatinase-associated lipocalin (NGAL) as a biomarker for acute renal injury after cardiac surgery. *Lancet* 2005; **365**: 1231–1238.
52. Mishra J, Mori K, Ma Q *et al.* Neutrophil gelatinase-associated lipocalin: a novel early urinary biomarker for cisplatin nephrotoxicity. *Am J Nephrol* 2004; **24**: 307–315.
53. Mishra J, Ma Q, Prada A *et al.* Identification of neutrophil gelatinase-associated lipocalin as a novel early urinary biomarker for ischemic renal injury. *J Am Soc Nephrol* 2003; **14**: 2534–2543.
54. Nickolas TL, O'Rourke MJ, Yang J *et al.* Sensitivity and specificity of a single emergency department measurement of urinary neutrophil gelatinase-associated lipocalin for diagnosing acute kidney injury. *Ann Intern Med* 2008; **148**: 810–819.
55. Constantin JM, Futier E, Perbet S *et al.* Plasma neutrophil gelatinase-associated lipocalin is an early marker of acute kidney injury in adult critically ill patients: a prospective study. *J Crit Care* 2010; **25**: 176.e1–176.e6.
56. Cruz DN, de Cal M, Garzotto F *et al.* Plasma neutrophil gelatinase-associated lipocalin is an early biomarker for acute kidney injury in an adult ICU population. *Intensive Care Med* 2009; **36**: 444–451.
57. Haase-Fielitz A, Bellomo R, Devarajan P *et al.* Novel and conventional serum biomarkers predicting acute kidney injury in adult cardiac surgery—a prospective cohort study. *Crit Care Med* 2009; **37**: 553–560.
58. Dent CL, Ma Q, Dastrala S *et al.* Plasma neutrophil gelatinase-associated lipocalin predicts acute kidney injury, morbidity and mortality after pediatric cardiac surgery: a prospective uncontrolled cohort study. *Crit Care* 2007; **11**: R127.
59. Malyszko J, Bachorzewska-Gajewska H, Sitniewska E *et al.* Serum neutrophil gelatinase-associated lipocalin as a marker of renal function in non-diabetic patients with stage 2–4 chronic kidney disease. *Ren Fail* 2008; **30**: 625–628.
60. Ko GJ, Grigoryev DN, Linfert D *et al.* Transcriptional analysis of kidneys during repair from AKI reveals possible roles for NGAL and KIM-1 as biomarkers of AKI to CKD transition. *Am J Physiol Renal Physiol* 2010; **298**: F1472–F1483.
61. Aghel A, Shrestha K, Mullens W *et al.* Serum neutrophil gelatinase-associated lipocalin (NGAL) in predicting worsening renal function in acute decompensated heart failure. *J Card Fail* 2010; **16**: 49–54.
62. Viau A, El Karoui K, Laouari D *et al.* Lipocalin 2 is essential for chronic kidney disease progression in mice and humans. *J Clin Invest* 2010; **120**: 4065–4076.
63. Malyszko J, Malyszko JS, Bachorzewska-Gajewska H *et al.* Neutrophil gelatinase-associated lipocalin is a new and sensitive marker of kidney function in chronic kidney disease patients and renal allograft recipients. *Transplant Proc* 2009; **41**: 158–161.
64. Ding H, He Y, Li K *et al.* Urinary neutrophil gelatinase-associated lipocalin (NGAL) is an early biomarker for renal tubulointerstitial injury in IgA nephropathy. *Clin Immunol* 2007; **123**: 227–234.
65. Bolignano D, Coppolino G, Campo S *et al.* Neutrophil gelatinase-associated lipocalin in patients with autosomal-dominant polycystic kidney disease. *Am J Nephrol* 2007; **27**: 373–378.
66. Bolignano D, Coppolino G, Campo S *et al.* Urinary neutrophil gelatinase-associated lipocalin (NGAL) is associated with severity of renal disease in proteinuric patients. *Nephrol Dial Transplant* 2008; **23**: 414–416.
67. Brunner HI, Mueller M, Rutherford C *et al.* Urinary neutrophil gelatinase-associated lipocalin as a biomarker of nephritis in childhood-onset systemic lupus erythematosus. *Arthritis Rheum* 2006; **54**: 2577–2584.
68. Mitsnefes MM, Kathman TS, Mishra J *et al.* Serum neutrophil gelatinase-associated lipocalin as a marker of renal function in children with chronic kidney disease. *Pediatr Nephrol* 2007; **22**: 101–108.
69. Bolignano D, Lacquaniti A, Coppolino G *et al.* Neutrophil gelatinase-associated lipocalin (NGAL) and progression of chronic kidney disease. *Clin J Am Soc Nephrol* 2009; **4**: 337–344.
70. Bolignano D, Lacquaniti A, Coppolino G *et al.* Neutrophil gelatinase-associated lipocalin as an early biomarker of nephropathy in diabetic patients. *Kidney Blood Press Res* 2009; **32**: 91–98.
71. Nishida M, Kawakatsu H, Okumura Y *et al.* Serum and urinary NGAL levels in children with chronic renal diseases. *Pediatr Int* 2010; **52**: 563–568.
72. Wu Y, Su T, Yang L *et al.* Urinary neutrophil gelatinase-associated lipocalin: a potential biomarker for predicting rapid progression of drug-induced chronic tubulointerstitial nephritis. *Am J Med Sci* 2010; **339**: 537–542.
73. Bolignano D, Donato V, Coppolino G *et al.* Neutrophil gelatinase-associated lipocalin (NGAL) as a marker of kidney damage. *Am J Kidney Dis* 2008; **52**: 595–605.
74. van Timmeren MM, van den Heuvel MC, Bailly V *et al.* Tubular kidney injury molecule-1 (KIM-1) in human renal disease. *J Pathol* 2007; **212**: 209–217.
75. Han WK, Bailly V, Abichandani R *et al.* Kidney injury molecule-1 (KIM-1): a novel biomarker for human renal proximal tubule injury. *Kidney Int* 2002; **62**: 237–244.
76. van Timmeren MM, Vaidya VS, van Ree RM *et al.* High urinary excretion of kidney injury molecule-1 is an independent predictor of graft loss in renal transplant recipients. *Transplantation* 2007; **84**: 1625–1630.
77. Waanders F, Vaidya VS, van Goor H *et al.* Effect of renin-angiotensin-aldosterone system inhibition, dietary sodium restriction, and/or diuretics on urinary kidney injury molecule 1 excretion in nondiabetic proteinuric kidney disease: a post hoc analysis of a randomized controlled trial. *Am J Kidney Dis* 2009; **53**: 16–25.
78. Vaidya VS, Ozer JS, Dieterle F *et al.* Kidney injury molecule-1 outperforms traditional biomarkers of kidney injury in preclinical biomarker qualification studies. *Nat Biotechnol* 2010; **28**: 478–485.
79. Vaidya VS, Niewczas MA, Ficociello LH *et al.* Regression of microalbuminuria in type 1 diabetes is associated with lower levels of urinary tubular injury biomarkers, kidney injury molecule-1, and N-acetyl-beta-D-glucosaminidase. *Kidney Int* 2011; **79**: 464–470.
80. Bosomworth MP, Aparicio SR, Hay AW. Urine N-acetyl-beta-D-glucosaminidase—a marker of tubular damage? *Nephrol Dial Transplant* 1999; **14**: 620–626.
81. Ellis EN, Brouhard BH, LaGrone L. Urinary N-acetyl-beta-D-glucosaminidase in streptozotocin-induced diabetic rats. *Biochem Med* 1984; **31**: 303–310.
82. Guder WG, Hofmann W. Markers for the diagnosis and monitoring of renal tubular lesions. *Clin Nephrol* 1992; **38**(Suppl 1): S3–S7.

83. Watts GF, Vlitos MA, Morris RW *et al.* Urinary N-acetyl-beta-D-glucosaminidase excretion in insulin-dependent diabetes mellitus: relation to microalbuminuria, retinopathy and glycaemic control. *Diabetes Metab* 1988; **14**: 653–658.
84. Kern EF, Erhard P, Sun W *et al.* Early urinary markers of diabetic kidney disease: a nested case-control study from the Diabetes Control and Complications Trial (DCCT). *Am J Kidney Dis* 2010; **55**: 824–834.
85. Holdt-Lehmann B, Lehmann A, Korten G *et al.* Diagnostic value of urinary alanine aminopeptidase and N-acetyl-beta-D-glucosaminidase in comparison to alpha 1-microglobulin as a marker in evaluating tubular dysfunction in glomerulonephritis patients. *Clin Chim Acta* 2000; **297**: 93–102.
86. Nakamura T, Sugaya T, Kawagoe Y *et al.* Effect of pitavastatin on urinary liver-type fatty acid-binding protein levels in patients with early diabetic nephropathy. *Diabetes Care* 2005; **28**: 2728–2732.
87. Ishimitsu T, Ohta S, Saito M *et al.* Urinary excretion of liver fatty acid-binding protein in health-check participants. *Clin Exp Nephrol* 2005; **9**: 34–39.
88. Kamijo A, Sugaya T, Hikawa A *et al.* Urinary liver-type fatty acid binding protein as a useful biomarker in chronic kidney disease. *Mol Cell Biochem* 2006; **284**: 175–182.
89. Kamijo A, Kimura K, Sugaya T *et al.* Urinary fatty acid-binding protein as a new clinical marker of the progression of chronic renal disease. *J Lab Clin Med* 2004; **143**: 23–30.
90. Kamijo A, Sugaya T, Hikawa A *et al.* Urinary excretion of fatty acid-binding protein reflects stress overload on the proximal tubules. *Am J Pathol* 2004; **165**: 1243–1255.
91. Kamijo A, Sugaya T, Hikawa A *et al.* Clinical evaluation of urinary excretion of liver-type fatty acid-binding protein as a marker for the monitoring of chronic kidney disease: a multicenter trial. *J Lab Clin Med* 2005; **145**: 125–133.
92. Nielsen SE, Sugaya T, Hovind P *et al.* Urinary liver-type fatty acid-binding protein (u-LFABP) predicts progression to nephropathy in type 1 diabetic patients. *Diabetes Care* 2010; **33**: 1320–1324.
93. Nakamura T, Sugaya T, Kawagoe Y *et al.* Effect of pitavastatin on urinary liver-type fatty-acid-binding protein in patients with nondiabetic mild chronic kidney disease. *Am J Nephrol* 2006; **26**: 82–86.
94. Truong LD, Foster SV, Barrios R *et al.* Tenascin is an ubiquitous extracellular matrix protein of human renal interstitium in normal and pathologic conditions. *Nephron* 1996; **72**: 579–586.
95. Jones CL, Buch S, Post M *et al.* Renal extracellular matrix accumulation in acute puromycin aminonucleoside nephrosis in rats. *Am J Pathol* 1992; **141**: 1381–1396.
96. Horstrup JH, Gehrmann M, Schneider B *et al.* Elevation of serum and urine levels of TIMP-1 and tenascin in patients with renal disease. *Nephrol Dial Transplant* 2002; **17**: 1005–1013.
97. Wang G, Lai FM, Lai KB *et al.* Messenger RNA expression of podocyte-associated molecules in the urinary sediment of patients with diabetic nephropathy. *Nephron Clin Pract* 2007; **106**: c169–c179.
98. Wang G, Lai FM, Tam LS *et al.* Messenger RNA expression of podocyte-associated molecules in urinary sediment of patients with lupus nephritis. *J Rheumatol* 2007; **34**: 2358–2364.
99. Kanno K, Kawachi H, Uchida Y *et al.* Urinary sediment podocalyxin in children with glomerular diseases. *Nephron Clin Pract* 2003; **95**: c91–c99.
100. Lajer M, Tarnow L, Jorsal A *et al.* Plasma concentration of asymmetric dimethylarginine (ADMA) predicts cardiovascular morbidity and mortality in type 1 diabetic patients with diabetic nephropathy. *Diabetes Care* 2008; **31**: 747–752.
101. Ravani P, Tripepi G, Malberti F *et al.* Asymmetrical dimethylarginine predicts progression to dialysis and death in patients with chronic kidney disease: a competing risks modeling approach. *J Am Soc Nephrol* 2005; **16**: 2449–2455.
102. Hanai K, Babazono T, Nyumura I *et al.* Asymmetric dimethylarginine is closely associated with the development and progression of nephropathy in patients with type 2 diabetes. *Nephrol Dial Transplant* 2009; **24**: 1884–1888.
103. Tsioufis C, Dimitriadis K, Andrikou E *et al.* ADMA, C-reactive protein, and albuminuria in untreated essential hypertension: a cross-sectional study. *Am J Kidney Dis* 2010; **55**: 1050–1059.
104. Vesely DL. Atrial natriuretic peptides in pathophysiological diseases. *Cardiovasc Res* 2001; **51**: 647–658.
105. Bunton DC, Petrie MC, Hillier C *et al.* The clinical relevance of adrenomedullin: a promising profile? *Pharmacol Ther* 2004; **103**: 179–201.
106. Epshteyn V, Morrison K, Krishnaswamy P *et al.* Utility of B-type natriuretic peptide (BNP) as a screen for left ventricular dysfunction in patients with diabetes. *Diabetes Care* 2003; **26**: 2081–2087.
107. Maisel AS, Krishnaswamy P, Nowak RM *et al.* Rapid measurement of B-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med* 2002; **347**: 161–167.
108. McCullough PA, Nowak RM, McCord J *et al.* B-type natriuretic peptide and clinical judgment in emergency diagnosis of heart failure: analysis from Breathing Not Properly (BNP) Multinational Study. *Circulation* 2002; **106**: 416–422.
109. Zaphiriou A, Robb S, Murray-Thomas T *et al.* The diagnostic accuracy of plasma BNP and NTproBNP in patients referred from primary care with suspected heart failure: results of the UK natriuretic peptide study. *Eur J Heart Fail* 2005; **7**: 537–541.
110. Gardner RS, Ozalp F, Murday AJ *et al.* N-terminal pro-brain natriuretic peptide. A new gold standard in predicting mortality in patients with advanced heart failure. *Eur Heart J* 2003; **24**: 1735–1743.
111. McDonagh TA, Robb SD, Murdoch DR *et al.* Biochemical detection of left-ventricular systolic dysfunction. *Lancet* 1998; **351**: 9–13.
112. Garg AX, Blake PG, Clark WF *et al.* Association between renal insufficiency and malnutrition in older adults: results from the NHANES III. *Kidney Int* 2001; **60**: 1867–1874.
113. Shlipak MG, Fried LF, Crump C *et al.* Elevations of inflammatory and procoagulant biomarkers in elderly persons with renal insufficiency. *Circulation* 2003; **107**: 87–92.
114. Pereira BJ, Shapiro L, King AJ *et al.* Plasma levels of IL-1 beta, TNF alpha and their specific inhibitors in undialyzed chronic renal failure, CAPD and hemodialysis patients. *Kidney Int* 1994; **45**: 890–896.
115. Knight EL, Rimm EB, Pai JK *et al.* Kidney dysfunction, inflammation, and coronary events: a prospective study. *J Am Soc Nephrol* 2004; **15**: 1897–1903.
116. Tonelli M, Sacks F, Pfeffer M *et al.* Biomarkers of inflammation and progression of chronic kidney disease. *Kidney Int* 2005; **68**: 237–245.
117. Fox ER, Benjamin EJ, Sarpong DF *et al.* The relation of C-reactive protein to chronic kidney disease in African Americans: the Jackson Heart Study. *BMC Nephrol* 2010; **11**: 1.
118. Fakhrazadeh H, Ghaderpanahi M, Sharifi F *et al.* Increased risk of chronic kidney disease in elderly with metabolic syndrome and high levels of C-reactive protein: Kahrizak Elderly Study. *Kidney Blood Press Res* 2009; **32**: 457–463.
119. Hung AM, Crawford DC, Griffin MR *et al.* CRP polymorphisms and progression of chronic kidney disease in African Americans. *Clin J Am Soc Nephrol* 2010; **5**: 24–33.
120. Alles VV, Bottazzi B, Peri G *et al.* Inducible expression of PTX3, a new member of the pentraxin family, in human mononuclear phagocytes. *Blood* 1994; **84**: 3483–3493.
121. Tong M, Carrero JJ, Qureshi AR *et al.* Plasma pentraxin 3 in patients with chronic kidney disease: associations with renal function, protein-energy wasting, cardiovascular disease, and mortality. *Clin J Am Soc Nephrol* 2007; **2**: 889–897.
122. Parikh CR, Jani A, Melnikov VY *et al.* Urinary interleukin-18 is a marker of human acute tubular necrosis. *Am J Kidney Dis* 2004; **43**: 405–414.
123. Nickolas TL, Barasch J, Devarajan P. Biomarkers in acute and chronic kidney disease. *Curr Opin Nephrol Hypertens* 2008; **17**: 127–132.
124. Miyauchi K, Takiyama Y, Honjyo J *et al.* Upregulated IL-18 expression in type 2 diabetic subjects with nephropathy: TGF-beta1 enhanced IL-18 expression in human renal proximal tubular epithelial cells. *Diabetes Res Clin Pract* 2009; **83**: 190–199.
125. Basile DP. The transforming growth factor beta system in kidney disease and repair: recent progress and future directions. *Curr Opin Nephrol Hypertens* 1999; **8**: 21–30.
126. Honkanen E, Teppo AM, Tornroth T *et al.* Urinary transforming growth factor-beta 1 in membranous glomerulonephritis. *Nephrol Dial Transplant* 1997; **12**: 2562–2568.
127. Fagerudd JA, Groop PH, Honkanen E *et al.* Urinary excretion of TGF-beta 1, PDGF-BB and fibronectin in insulin-dependent diabetes mellitus patients. *Kidney Int Suppl* 1997; **63**: S195–S197.
128. Suthanthiran M, Gerber LM, Schwartz JE *et al.* Circulating transforming growth factor-beta1 levels and the risk for kidney disease in African Americans. *Kidney Int* 2009; **76**: 72–80.
129. Gilbert RE, Akdeniz A, Weitz S *et al.* Urinary connective tissue growth factor excretion in patients with type 1 diabetes and nephropathy. *Diabetes Care* 2003; **26**: 2632–2636.
130. Vanden Heuvel GB. CD14: a candidate biomarker for the prognosis of polycystic kidney disease. *Kidney Int* 2010; **78**: 537–538.

131. Zhou J, Ouyang X, Cui X *et al.* Renal CD14 expression correlates with the progression of cystic kidney disease. *Kidney Int* 2010; **78**: 550–560.
132. Liu S, Tang W, Zhou J *et al.* Fibroblast growth factor 23 is a counter-regulatory phosphaturic hormone for vitamin D. *J Am Soc Nephrol* 2006; **17**: 1305–1315.
133. Westerberg PA, Linde T, Wikstrom B *et al.* Regulation of fibroblast growth factor-23 in chronic kidney disease. *Nephrol Dial Transplant* 2007; **22**: 3202–3207.
134. Kazama JJ, Sato F, Omori K *et al.* Pretreatment serum FGF-23 levels predict the efficacy of calcitriol therapy in dialysis patients. *Kidney Int* 2005; **67**: 1120–1125.
135. Juppner H. Novel regulators of phosphate homeostasis and bone metabolism. *Ther Apher Dial* 2007; **11**(Suppl 1): S3–S22.
136. Gutierrez O, Isakova T, Rhee E *et al.* Fibroblast growth factor-23 mitigates hyperphosphatemia but accentuates calcitriol deficiency in chronic kidney disease. *J Am Soc Nephrol* 2005; **16**: 2205–2215.
137. Manghat P, Fraser WD, Wierzbicki AS *et al.* Fibroblast growth factor-23 is associated with C-reactive protein, serum phosphate and bone mineral density in chronic kidney disease. *Osteoporos Int* 2010; **21**: 1853–1861.
138. Oliveira RB, Cancela AL, Gracioli FG *et al.* Early control of PTH and FGF23 in normophosphatemic CKD patients: a new target in CKD-MBD therapy? *Clin J Am Soc Nephrol* 2010; **5**: 286–291.
139. Kronenberg F, Konig P, Neyer U *et al.* Multicenter study of lipoprotein(a) and apolipoprotein(a) phenotypes in patients with end-stage renal disease treated by hemodialysis or continuous ambulatory peritoneal dialysis. *J Am Soc Nephrol* 1995; **6**: 110–120.
140. Kronenberg F, Kuen E, Ritz E *et al.* Apolipoprotein A-IV serum concentrations are elevated in patients with mild and moderate renal failure. *J Am Soc Nephrol* 2002; **13**: 461–469.
141. Scherer PE, Williams S, Fogliano M *et al.* A novel serum protein similar to C1q, produced exclusively in adipocytes. *J Biol Chem* 1995; **270**: 26746–26749.
142. Zoccali C, Mallamaci F, Panuccio V *et al.* Adiponectin is markedly increased in patients with nephrotic syndrome and is related to metabolic risk factors. *Kidney Int Suppl* 2003; S98–S102.
143. Stenvinkel P, Marchlewska A, Pecoits-Filho R *et al.* Adiponectin in renal disease: relationship to phenotype and genetic variation in the gene encoding adiponectin. *Kidney Int* 2004; **65**: 274–281.
144. Saraheimo M, Forsblom C, Thorn L *et al.* Serum adiponectin and progression of diabetic nephropathy in patients with type 1 diabetes. *Diabetes Care* 2008; **31**: 1165–1169.
145. Holvoet P, Donck J, Landeloos M *et al.* Correlation between oxidized low density lipoproteins and von Willebrand factor in chronic renal failure. *Thromb Haemost* 1996; **76**: 663–669.
146. Witko-Sarsat V, Friedlander M, Capeillere-Blandin C *et al.* Advanced oxidation protein products as a novel marker of oxidative stress in uremia. *Kidney Int* 1996; **49**: 1304–1313.
147. Oberg BP, McMenamin E, Lucas FL *et al.* Increased prevalence of oxidant stress and inflammation in patients with moderate to severe chronic kidney disease. *Kidney Int* 2004; **65**: 1009–1016.
148. Mezzano D, Pais EO, Aranda E *et al.* Inflammation, not hyperhomocysteinemia, is related to oxidative stress and hemostatic and endothelial dysfunction in uremia. *Kidney Int* 2001; **60**: 1844–1850.
149. Dincer Y, Sekercioglu N, Pekpak M *et al.* Assessment of DNA oxidation and antioxidant activity in hypertensive patients with chronic kidney disease. *Ren Fail* 2008; **30**: 1006–1011.
150. Dounousi E, Papavasiliou E, Makedou A *et al.* Oxidative stress is progressively enhanced with advancing stages of CKD. *Am J Kidney Dis* 2006; **48**: 752–760.
151. Calabrese V, Mancuso C, Sapienza M *et al.* Oxidative stress and cellular stress response in diabetic nephropathy. *Cell Stress Chaperones* 2007; **12**: 299–306.
152. Beisswenger PJ, Drummond KS, Nelson RG *et al.* Susceptibility to diabetic nephropathy is related to dicarbonyl and oxidative stress. *Diabetes* 2005; **54**: 3274–3281.
153. Gugliucci A, Bendayan M. Renal fate of circulating advanced glycosylated end products (AGE): evidence for reabsorption and catabolism of AGE-peptides by renal proximal tubular cells. *Diabetologia* 1996; **39**: 149–160.
154. Simmons EM, Langone A, Sezer MT *et al.* Effect of renal transplantation on biomarkers of inflammation and oxidative stress in end-stage renal disease patients. *Transplantation* 2005; **79**: 914–919.
155. Whitfield JB. Serum gamma-glutamyltransferase and risk of disease. *Clin Chem* 2007; **53**: 1–2.
156. Conigrave KM, Saunders JB, Reznik RB *et al.* Prediction of alcohol-related harm by laboratory test results. *Clin Chem* 1993; **39**(11 Part 1): 2266–2270.
157. Lee DH, Blomhoff R, Jacobs Jr DR. Is serum gamma glutamyltransferase a marker of oxidative stress? *Free Radic Res* 2004; **38**: 535–539.
158. Lee DH, Jacobs Jr DR, Gross M *et al.* Serum gamma-glutamyltransferase was differently associated with microalbuminuria by status of hypertension or diabetes: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Clin Chem* 2005; **51**: 1185–1191.
159. Ryu S, Chang Y, Kim DI *et al.* gamma-Glutamyltransferase as a predictor of chronic kidney disease in nonhypertensive and nondiabetic Korean men. *Clin Chem* 2007; **53**: 71–77.
160. Targher G, Kendrick J, Smits G *et al.* Relationship between serum gamma-glutamyltransferase and chronic kidney disease in the United States adult population. Findings from the National Health and Nutrition Examination Survey 2001–2006. *Nutr Metab Cardiovasc Dis* 2010; **20**: 583–590.
161. Teppala S, Shankar A, Li J *et al.* Association between serum gamma-glutamyltransferase and chronic kidney disease among US adults. *Kidney Blood Press Res* 2010; **33**: 1–6.
162. Beck Jr LH, Bonegio RG, Lambeau G *et al.* M-type phospholipase A2 receptor as target antigen in idiopathic membranous nephropathy. *N Engl J Med* 2009; **361**: 11–21.
163. Liu YH, Chen CH, Chen SY *et al.* Association of phospholipase A2 receptor 1 polymorphisms with idiopathic membranous nephropathy in Chinese patients in Taiwan. *J Biomed Sci* 2010; **17**: 81.
164. Kim S, Chin HJ, Na KY *et al.* Single nucleotide polymorphisms in the phospholipase A(2) receptor gene are associated with genetic susceptibility to idiopathic membranous nephropathy. *Nephron Clin Pract* 2011; **117**: c253–c258.
165. Taglieri N, Koenig W, Kaski JC. Cystatin C and cardiovascular risk. *Clin Chem* 2009; **55**: 1932–1943.
166. Shlipak MG, Katz R, Sarnak MJ *et al.* Cystatin C and prognosis for cardiovascular and kidney outcomes in elderly persons without chronic kidney disease. *Ann Intern Med* 2006; **145**: 237–246.
167. Rifkin DE, Katz R, Chonchol M *et al.* Albuminuria, impaired kidney function and cardiovascular outcomes or mortality in the elderly. *Nephrol Dial Transplant* 2010; **25**: 1560–1567.
168. Kiyosue A, Hirata Y, Ando J *et al.* Plasma cystatin C concentration reflects the severity of coronary artery disease in patients without chronic kidney disease. *Circ J* 2010; **74**: 2441–2447.
169. Ix JH, Shlipak MG, Chertow GM *et al.* Association of cystatin C with mortality, cardiovascular events, and incident heart failure among persons with coronary heart disease: data from the Heart and Soul Study. *Circulation* 2007; **115**: 173–179.
170. Navaneethan SD, Beddhu S. Associations of serum uric acid with cardiovascular events and mortality in moderate chronic kidney disease. *Nephrol Dial Transplant* 2009; **24**: 1260–1266.
171. Sakuma M, Nakamura M, Tanaka F *et al.* Plasma B-type natriuretic peptide level and cardiovascular events in chronic kidney disease in a community-based population. *Circ J* 2010; **74**: 792–797.
172. Hedvig J, Podracka L, Potoceková D. Elevated brain natriuretic peptide is associated with abnormal heart geometry in children with chronic kidney disease. *Kidney Blood Press Res* 2010; **33**: 87–93.
173. Cooke JP. Asymmetrical dimethylarginine: the Uber marker? *Circulation* 2004; **109**: 1813–1818.
174. Kielstein JT, Impraïm B, Simmel S *et al.* Cardiovascular effects of systemic nitric oxide synthase inhibition with asymmetrical dimethylarginine in humans. *Circulation* 2004; **109**: 172–177.
175. Stuhlinger MC, Abbasi F, Chu JW *et al.* Relationship between insulin resistance and an endogenous nitric oxide synthase inhibitor. *JAMA* 2002; **287**: 1420–1426.
176. Zoccali C, Bode-Boger S, Mallamaci F *et al.* Plasma concentration of asymmetrical dimethylarginine and mortality in patients with end-stage renal disease: a prospective study. *Lancet* 2001; **358**: 2113–2117.
177. Wilhelmsen L, Svardsudd K, Korsan-Bengtson K *et al.* Fibrinogen as a risk factor for stroke and myocardial infarction. *N Engl J Med* 1984; **311**: 501–505.
178. Kannel WB, Wolf PA, Castelli WP *et al.* Fibrinogen and risk of cardiovascular disease. The Framingham Study. *JAMA* 1987; **258**: 1183–1186.

179. Ernst E, Resch KL. Fibrinogen as a cardiovascular risk factor: a meta-analysis and review of the literature. *Ann Intern Med* 1993; **118**: 956–963.
180. Danesh J, Collins R, Appleby P *et al.* Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA* 1998; **279**: 1477–1482.
181. Zoccali C, Mallamaci F, Tripepi G *et al.* Fibrinogen, mortality and incident cardiovascular complications in end-stage renal failure. *J Intern Med* 2003; **254**: 132–139.
182. Shlipak MG, Fried LF, Cushman M *et al.* Cardiovascular mortality risk in chronic kidney disease: comparison of traditional and novel risk factors. *JAMA* 2005; **293**: 1737–1745.
183. Muntner P, He J, Astor BC *et al.* Traditional and nontraditional risk factors predict coronary heart disease in chronic kidney disease: results from the atherosclerosis risk in communities study. *J Am Soc Nephrol* 2005; **16**: 529–538.
184. Goicoechea M, de Vinuesa SG, Gomez-Campdera F *et al.* Serum fibrinogen levels are an independent predictor of mortality in patients with chronic kidney disease (CKD) stages 3 and 4. *Kidney Int Suppl* 2008; **111**: S67–S70.
185. Weiner DE, Tighiouart H, Elsayed EF *et al.* The relationship between nontraditional risk factors and outcomes in individuals with stage 3 to 4 CKD. *Am J Kidney Dis* 2008; **51**: 212–223.
186. Bolignano D, Coppolino G, Lacquaniti A *et al.* From kidney to cardiovascular diseases: NGAL as a biomarker beyond the confines of nephrology. *Eur J Clin Invest* 2010; **40**: 273–276.
187. Bu DX, Hemdahl AL, Gabrielsen A *et al.* Induction of neutrophil gelatinase-associated lipocalin in vascular injury via activation of nuclear factor-kappaB. *Am J Pathol* 2006; **169**: 2245–2253.
188. Folkesson M, Kazi M, Zhu C *et al.* Presence of NGAL/MMP-9 complexes in human abdominal aortic aneurysms. *Thromb Haemost* 2007; **98**: 427–433.
189. Elneihoum AM, Falke P, Hedblad B *et al.* Leukocyte activation in atherosclerosis: correlation with risk factors. *Atherosclerosis* 1997; **131**: 79–84.
190. Bolignano D, Basile G, Parisi P *et al.* Increased plasma neutrophil gelatinase-associated lipocalin levels predict mortality in elderly patients with chronic heart failure. *Rejuvenation Res* 2009; **12**: 7–14.
191. Seiler S, Reichart B, Roth D *et al.* FGF-23 and future cardiovascular events in patients with chronic kidney disease before initiation of dialysis treatment. *Nephrol Dial Transplant* 2010; **25**: 3983–3989.
192. Kanbay M, Nicoleta M, Selcoki Y *et al.* Fibroblast growth factor 23 and fetuin A are independent predictors for the coronary artery disease extent in mild chronic kidney disease. *Clin J Am Soc Nephrol* 2010; **5**: 1780–1786.
193. Wolf M. FGF23: fashion or physiology? *Clin J Am Soc Nephrol* 2010; **5**: 1727–1729.
194. Wiley SR, Schooley K, Smolak PJ *et al.* Identification and characterization of a new member of the TNF family that induces apoptosis. *Immunity* 1995; **3**: 673–682.
195. Michowitz Y, Goldstein E, Roth A *et al.* The involvement of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) in atherosclerosis. *J Am Coll Cardiol* 2005; **45**: 1018–1024.
196. Liabeuf S, Barreto DV, Barreto FC *et al.* The circulating soluble TRAIL is a negative marker for inflammation inversely associated with the mortality risk in chronic kidney disease patients. *Nephrol Dial Transplant* 2010; **25**: 2596–2602.
197. Apple FS, Murakami MM, Pearce LA *et al.* Multi-biomarker risk stratification of N-terminal pro-B-type natriuretic peptide, high-sensitivity C-reactive protein, and cardiac troponin T and I in end-stage renal disease for all-cause death. *Clin Chem* 2004; **50**: 2279–2285.
198. Kiatchoosakun S, Sahasthas D, Wongvipaporn C *et al.* Cardiac troponin-T in pre-end stage kidney disease. *J Med Assoc Thai* 2008; **91**: 1806–1811.
199. Tsutamoto T, Kawahara C, Yamaji M *et al.* Relationship between renal function and serum cardiac troponin T in patients with chronic heart failure. *Eur J Heart Fail* 2009; **11**: 653–658.
200. Abbas NA, John RI, Webb MC *et al.* Cardiac troponins and renal function in nondialysis patients with chronic kidney disease. *Clin Chem* 2005; **51**: 2059–2066.
201. Lamb EJ, Kenny C, Abbas NA *et al.* Cardiac troponin I concentration is commonly increased in nondialysis patients with CKD: experience with a sensitive assay. *Am J Kidney Dis* 2007; **49**: 507–516.
202. Hayashi T, Obi Y, Kimura T *et al.* Cardiac troponin T predicts occult coronary artery stenosis in patients with chronic kidney disease at the start of renal replacement therapy. *Nephrol Dial Transplant* 2008; **23**: 2936–2942.
203. Melloni C, Alexander KP, Milford-Beland S *et al.* Prognostic value of troponins in patients with non-ST-segment elevation acute coronary syndromes and chronic kidney disease. *Clin Cardiol* 2008; **31**: 125–129.
204. Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med* 2005; **352**: 1685–1695.
205. Frankenberger M, Sternsdorf T, Pechumer H *et al.* Differential cytokine expression in human blood monocyte subpopulations: a polymerase chain reaction analysis. *Blood* 1996; **87**: 373–377.
206. Heine GH, Ulrich C, Seibert E *et al.* CD14(++)CD16+ monocytes but not total monocyte numbers predict cardiovascular events in dialysis patients. *Kidney Int* 2008; **73**: 622–629.
207. Rogacev KS, Ulrich C, Blomer L *et al.* Monocyte heterogeneity in obesity and subclinical atherosclerosis. *Eur Heart J* 2010; **31**: 369–376.
208. Ozer JS, Dieterle F, Troth S *et al.* A panel of urinary biomarkers to monitor reversibility of renal injury and a serum marker with improved potential to assess renal function. *Nat Biotechnol* 2010; **28**: 486–494.
209. Dieterle F, Perentes E, Cordier A *et al.* Urinary clusterin, cystatin C, beta2-microglobulin and total protein as markers to detect drug-induced kidney injury. *Nat Biotechnol* 2010; **28**: 463–469.
210. Yu Y, Jin H, Holder D *et al.* Urinary biomarkers trefoil factor 3 and albumin enable early detection of kidney tubular injury. *Nat Biotechnol* 2010; **28**: 470–477.
211. Dieterle F, Sistare F, Goodsaid F *et al.* Renal biomarker qualification submission: a dialog between the FDA-EMEA and Predictive Safety Testing Consortium. *Nat Biotechnol* 2010; **28**: 455–462.
212. Levin A, Djurdjev O, Beaulieu M *et al.* Variability and risk factors for kidney disease progression and death following attainment of stage 4 CKD in a referred cohort. *Am J Kidney Dis* 2008; **52**: 661–671.
213. Sayers S, Singh G, Mott S *et al.* Relationships between birthweight and biomarkers of chronic disease in childhood: Aboriginal Birth Cohort Study 1987–2001. *Paediatr Perinat Epidemiol* 2009; **23**: 548–556.
214. Keller C, Katz R, Sarnak MJ *et al.* Inflammatory biomarkers and decline in kidney function in the elderly: the Cardiovascular Health Study. *Nephrol Dial Transplant* 2010; **25**: 119–124.